

## CIRCULAR ORBITS: RELATION BETWEEN RADIUS AND ANGULAR MOMENTUM

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 10; Problems 10.7-8.

For a circular orbit, the relation between the angular momentum and radius is

$$(1) \quad r = \frac{6GM}{1 \mp \sqrt{1 - 12(GM/l)^2}}$$

We can invert this to get  $l$  in terms of  $r$ :

$$(2) \quad \mp \sqrt{1 - 12(GM/l)^2} = \frac{6GM}{r} - 1$$

$$(3) \quad -12 \frac{G^2 M^2}{l^2} = \left( \frac{6GM}{r} - 1 \right)^2 - 1$$

$$(4) \quad l^2 = \frac{12r^2 (GM)^2}{r^2 - (6GM - r)^2}$$

$$(5) \quad = \frac{r^2 GM}{r - 3GM}$$

From this we can get an expression for  $\tilde{E}$  for a circular orbit where  $dr/d\tau = 0$ :

$$\begin{aligned}
 (6) \quad \tilde{E} &= \frac{1}{2} \frac{l^2}{r^2} - GM \left( \frac{1}{r} + \frac{l^2}{r^3} \right) \\
 (7) \quad &= -\frac{GM}{r} + \frac{r^2 GM}{r - 3GM} \left( \frac{1}{2r^2} - \frac{GM}{r^3} \right) \\
 (8) \quad &= -\frac{GM}{2r} \left[ 2 - \frac{2r^3}{r - 3GM} \left( \frac{1}{2r^2} - \frac{GM}{r^3} \right) \right] \\
 (9) \quad &= -\frac{GM}{2r} \left[ \frac{r}{r - 3GM} \left( \frac{2r - 6GM}{r} - \left( 1 - \frac{2GM}{r} \right) \right) \right] \\
 (10) \quad &= -\frac{GM}{2r} \left( 1 - \frac{3GM}{r} \right)^{-1} \left( 1 - \frac{4GM}{r} \right)
 \end{aligned}$$

The energy per unit mass  $e$  is then, using  $\tilde{E} = \frac{1}{2}(e^2 - 1)$ :

$$(11) \quad e = \left[ 1 - \frac{GM}{r} \left( 1 - \frac{3GM}{r} \right)^{-1} \left( 1 - \frac{4GM}{r} \right) \right]^{1/2}$$

For  $r = 6GM$ , this comes out to  $e = \sqrt{\frac{8}{9}}$ .

As an example of the use of these formulas, suppose we start an object at infinity with no radial velocity, but with an infinitesimal tangential velocity which gives it an angular momentum  $l$ . Since the object is at an infinite distance, the formula  $l = r^2 \omega$  means that in the limit as  $r \rightarrow \infty$ ,  $\omega \rightarrow 0$  so that the product  $r^2 \omega = l$ , thus the tangential motion really *is* infinitesimal.

As the object falls in towards the central mass, it spirals in, keeping  $l$  constant. Since the object started off essentially at rest,  $e = 1$  so one solution is for the object to end up in a circular orbit with  $r = 4GM$ . From the above formula, this corresponds to

$$(12) \quad l = 4GM \sqrt{\frac{GM}{4GM - 3GM}}$$

$$(13) \quad = 4GM$$

#### PINGBACKS

Pingback: Circular orbit around a supermassive black hole

Pingback: Circular orbit: Schwarzschild vs Newton

Pingback: Orbit of a comet around a black hole

Pingback: Twin paradox with a black hole

Pingback: Perihelion shift in planetary orbits

CIRCULAR ORBITS: RELATION BETWEEN RADIUS AND ANGULAR MOMENTUM 3

Pingback: Circular orbit: appearance to a falling observer

Pingback: Local flat frame for a circular orbit

Pingback: Perihelion shift: numerical solution