

## FALLING INTO A BLACK HOLE

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 14; Problems 14.1, 14.2.

Here are a couple of simple examples of calculating proper time for an object crossing the event horizon. If the object starts at rest at  $r = R$  and then falls to  $r = r_0$ , the proper time interval is

$$(1) \quad \Delta\tau = \sqrt{\frac{R}{8GM}} \left[ 2\sqrt{r(R-r)} + R \arctan \left( \frac{R-2r}{2\sqrt{r(R-r)}} \right) \right] \Big|_{r_0}^R$$

An alternative form of this is found by using the identity:

$$(2) \quad \arctan x = \arcsin \frac{x}{\sqrt{1+x^2}}$$

A bit of algebra produces:

$$(4) \quad \left( \frac{R-2r}{2\sqrt{r(R-r)}} \right) \left( 1 + \left[ \frac{R-2r}{2\sqrt{r(R-r)}} \right]^2 \right)^{-1/2} = \left( \frac{R-2r}{2\sqrt{r(R-r)}} \right) \frac{2\sqrt{r(R-r)}}{R} = \frac{R-2r}{R}$$

Therefore

$$(5) \quad \Delta\tau = \sqrt{\frac{R}{8GM}} \left[ 2\sqrt{r(R-r)} + R \arcsin \left( \frac{R-2r}{R} \right) \right] \Big|_{r_0}^R$$

**Example 1:** If we start at radius of  $R = 10GM$  outside a black hole with a million solar masses, then

$$(6) \quad GM = 10^6 (GM)_{sun}$$

$$(7) \quad = 1.477 \times 10^6 \text{ km}$$

The time to fall to zero radius ( $r_0 = 0$ ) is then

$$\begin{aligned}
(8) \quad \Delta\tau &= \frac{\pi r_2^{3/2}}{\sqrt{8GM}} \\
(9) \quad &= \pi \sqrt{\frac{1000}{8}} GM \\
(10) \quad &= 5.188 \times 10^7 \text{ km} = 172.9 \text{ sec}
\end{aligned}$$

**Example 2:** We now start at a radius of  $r_1 = 16GM$  from the same black hole, and launch an object radially outwards with a speed such that it comes to rest at  $r_2 = 32GM$ . The object then falls radially inwards until it crosses the event horizon at  $r_3 = 2GM$  and then continues until it reaches  $r_4 = 0$ . We can work out the times for each leg of the journey separately. We can get these times by inserting the appropriate limits in 5.

For the first leg, the time is the same as if the object started at  $R = 32GM$  and fell to  $r_0 = 16GM$ . This comes out to

$$(11) \quad \Delta\tau_{1 \rightarrow 2} = 32(2 + \pi) GM = 2.43 \times 10^8 \text{ km} = 810 \text{ sec}$$

Next, the object falls from rest at  $R = 32GM$  to the event horizon at  $r_0 = 2GM$ :

$$(12) \quad \Delta\tau_{2 \rightarrow 3} = 8 \left( \sqrt{15} + 8 \arcsin\left(\frac{7}{8}\right) + 4\pi \right) GM$$

$$(13) \quad = 199.7 GM = 2.95 \times 10^8 \text{ km} = 983.2 \text{ sec}$$

Finally, the object falls from  $R = 2GM$  to  $r_0 = 0$  which takes time

$$(14) \quad \Delta\tau_{3 \rightarrow 4} = \pi GM = 15.47 \text{ sec}$$