

EVENT HORIZON: PROPER TIME TO FALL INTO $R = 0$

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 14; Box 14.2.

Continuing our exploration of the Schwarzschild metric and its behaviour at $r = 2GM$ (known as the event horizon), we'll look here at the proper time an object measures as it crosses the event horizon. The metric is:

$$ds^2 = - \left(1 - \frac{2GM}{r}\right) dt^2 + \left(1 - \frac{2GM}{r}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (1)$$

Consider an object starting at rest at a radial coordinate $r = R$ and falling radially in towards the event horizon. To work out its proper time, we start with the equation of motion for r :

$$\frac{1}{2} \left(\frac{dr}{d\tau}\right)^2 + \frac{1}{2} \frac{\ell^2}{r^2} - GM \left(\frac{1}{r} + \frac{\ell^2}{r^3}\right) = \frac{1}{2} (e^2 - 1) \quad (2)$$

Since the object is at rest initially, $dr/d\tau = 0$ and its angular momentum is $\ell = 0$ so we get for its energy

$$e = \sqrt{1 - \frac{2GM}{R}} \quad (3)$$

The energy e is a conserved quantity, so the equation of motion becomes

$$\frac{dr}{d\tau} = \sqrt{2GM} \sqrt{\frac{1}{r} - \frac{1}{R}} \quad (4)$$

We can integrate this to find the proper time that elapses as the object falls from $r = R$ to $r = 0$:

$$\int_R^0 \frac{-dr}{\sqrt{\frac{1}{r} - \frac{1}{R}}} = \sqrt{2GM} \Delta\tau \quad (5)$$

(the minus sign in the integrand accounts for the fact that r is decreasing).

Using Maple, the integral is

$$\int \frac{-dr}{\sqrt{\frac{1}{r} - \frac{1}{R}}} = \frac{\sqrt{R}}{2} \left[2\sqrt{r(R-r)} + R \arctan \left(\frac{R-2r}{2\sqrt{r(R-r)}} \right) \right] \quad (6)$$

Plugging in the limits, we get

$$\sqrt{2GM}\Delta\tau = \frac{\pi R^{3/2}}{2} \quad (7)$$

$$\Delta\tau = \frac{\pi R^{3/2}}{\sqrt{8GM}} \quad (8)$$

PINGBACKS

Pingback: Falling into a black hole

Pingback: Event horizon: time and space swap round

Pingback: Kruskal-Szekeles metric: what can you see as you fall into a black hole?

Pingback: Black hole radiation: energy of emitted particles

Pingback: Falling into a black hole: tidal forces