

## CHRISTOFFEL SYMBOLS: SYMMETRY

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 17; Box 17.3.

The Christoffel symbols are defined in terms of the basis vectors in a given coordinate system as:

$$(0.1) \quad \boxed{\frac{\partial \mathbf{e}_i}{\partial x^j} = \Gamma_{ij}^k \mathbf{e}_k}$$

Remember that the basis vectors  $\mathbf{e}_i$  are defined so that

$$(0.2) \quad ds^2 = ds \cdot ds$$

$$(0.3) \quad = (dx^i \mathbf{e}_i) \cdot (dx^j \mathbf{e}_j)$$

$$(0.4) \quad = \mathbf{e}_i \cdot \mathbf{e}_j dx^i dx^j$$

$$(0.5) \quad \equiv g_{ij} dx^i dx^j$$

In a locally flat frame using rectangular spatial coordinates, the basis vectors  $\mathbf{e}_i$  are all constants, so from 0.1, all the Christoffel symbols must be zero:  $\Gamma_{ij}^k = 0$ .

Now let's look at the second covariant derivative of a scalar field  $\Phi$ :

$$(0.6) \quad \nabla_i \nabla_j \Phi = \nabla_i (\partial_j \Phi)$$

$$(0.7) \quad = \partial_i \partial_j \Phi - \Gamma_{ij}^k \partial_k \Phi$$

where in 0.6 we used rule 1 for the covariant derivative: the covariant derivative of a scalar is the same as the ordinary derivative.

In the locally flat frame, this equation reduces to

$$(0.8) \quad \nabla_i \nabla_j \Phi = \partial_i \partial_j \Phi$$

Since the covariant derivative is a tensor, this is a tensor equation, and since ordinary partial derivatives commute, this equation is the same if we swap the indices  $i$  and  $j$ . Tensor equations must have the same form in all coordinate systems, so this implies that 0.7 must also be invariant if we

swap  $i$  and  $j$ . This means that the Christoffel symbols are symmetric under exchange of their two lower indices:

$$(0.9) \quad \boxed{\Gamma_{ij}^k = \Gamma_{ji}^k}$$

At first glance, this seems wrong, since from the definition 0.1 this symmetry implies that

$$(0.10) \quad \frac{\partial \mathbf{e}_i}{\partial x^j} = \frac{\partial \mathbf{e}_j}{\partial x^i}$$

In 2-D polar coordinates, if we take the usual unit vectors  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$  then both these vectors are constants as we change  $r$  and both of them change when we change  $\theta$ , so it's certainly not true that  $\partial \hat{\mathbf{r}} / \partial \theta = \partial \hat{\boldsymbol{\theta}} / \partial r$ , for example. However, remember that the basis vectors we're using are *not* the usual unit vectors; rather they are defined so that condition 0.4 is true. In polar coordinates, we have

$$(0.11) \quad ds^2 = dr^2 + r^2 d\theta^2$$

so

$$(0.12) \quad \mathbf{e}_r = \hat{\mathbf{r}} = \cos \theta \hat{\mathbf{x}} + \sin \theta \hat{\mathbf{y}}$$

$$(0.13) \quad \mathbf{e}_\theta = r \hat{\boldsymbol{\theta}} = -r \sin \theta \hat{\mathbf{x}} + r \cos \theta \hat{\mathbf{y}}$$

For the derivatives, we have

$$(0.14) \quad \frac{\partial \mathbf{e}_r}{\partial \theta} = -\sin \theta \hat{\mathbf{x}} + \cos \theta \hat{\mathbf{y}} = \hat{\boldsymbol{\theta}}$$

$$(0.15) \quad \frac{\partial \mathbf{e}_\theta}{\partial r} = -\sin \theta \hat{\mathbf{x}} + \cos \theta \hat{\mathbf{y}} = \hat{\boldsymbol{\theta}}$$

Thus the condition 0.10 is actually satisfied here.

#### PINGBACKS

Pingback: Christoffel symbols for Schwarzschild metric

Pingback: Christoffel symbols in terms of the metric tensor

Pingback: Gravitoelectric and gravitomagnetic densities