

## RIEMANN TENSOR IN 2-D POLAR COORDINATES

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 18; Problem P18.4.

As a simple example of the Riemann tensor we can work out one of its components in 2-d polar coordinates. The tensor is

$$(0.1) \quad R^i{}_{j\ell m} \equiv \partial_\ell \Gamma^i{}_{mj} - \partial_m \Gamma^i{}_{\ell j} + \Gamma^k{}_{mj} \Gamma^i{}_{\ell k} - \Gamma^k{}_{\ell j} \Gamma^i{}_{km}$$

As usual, we need the Christoffel symbols, which we can get by comparing the two forms of the geodesic equation. These equations are

$$(0.2) \quad g_{aj} \ddot{x}^j + \left( \partial_i g_{aj} - \frac{1}{2} \partial_a g_{ij} \right) \dot{x}^j \dot{x}^i = 0$$

$$(0.3) \quad \ddot{x}^m + \Gamma^m{}_{ij} \dot{x}^j \dot{x}^i = 0$$

The metric is

$$(0.4) \quad ds^2 = dr^2 + r^2 d\theta^2$$

so  $g_{rr} = 1$  and  $g_{\theta\theta} = r^2$ . For the two coordinates, 0.2 gives us

$$(0.5) \quad \ddot{r} - r\dot{\theta}^2 = 0$$

$$(0.6) \quad r^2 \ddot{\theta} + 2r\dot{r}\dot{\theta} = 0$$

Comparing with 0.3 we get

$$(0.7) \quad \Gamma^r{}_{\theta\theta} = -r$$

$$(0.8) \quad \Gamma^{\theta}{}_{r\theta} = \Gamma^{\theta}{}_{\theta r} = \frac{1}{r}$$

One component of the Riemann tensor is :

$$(0.9) \quad R^r_{\theta r \theta} = \partial_r \Gamma^r_{\theta \theta} - \partial_\theta \Gamma^r_{r \theta} + \Gamma^r_{kr} \Gamma^k_{\theta \theta} - \Gamma^r_{\theta k} \Gamma^k_{r \theta}$$

$$(0.10) \quad = -1 - 0 + 0 + 1$$

$$(0.11) \quad = 0$$

In fact, there is only one independent component of  $R$  in a 2-d metric (as we'll see when we study the symmetry of  $R$  later), so all other components can be written as multiples of  $R^r_{\theta r \theta}$ . Thus for polar coordinates  $R^r_{\theta r \theta} = 0$  implies all components of  $R$  are zero, which is what we'd expect since polar coordinates describe flat space.