

RIEMANN TENSOR FOR 3-D SPHERICAL COORDINATES

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 19; Problem 19.7.

We saw that the Riemann tensor for the surface of a sphere had a non-zero component, indicating that this is a curved space. If we use spherical coordinates in 3-d space, however, the Riemann tensor should be zero, since this is a flat space.

As usual, we need the Christoffel symbols and we get them by comparing the two forms of the geodesic equation.

$$(0.1) \quad \frac{d}{d\tau} (g_{aj}\dot{x}^j) - \frac{1}{2}\partial_a g_{ij}\dot{x}^i\dot{x}^j = 0$$

$$(0.2) \quad \ddot{x}^m + \Gamma_{ij}^m\dot{x}^j\dot{x}^i = 0$$

where as usual a dot denotes a derivative with respect to proper time τ .

For spherical coordinates, the interval is

$$(0.3) \quad ds^2 = dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$$

Note that r is now a variable, rather than the constant radius of the sphere in the 2-d system.

From 0.1 we get, with $a = \theta$:

$$(0.4) \quad 2r\dot{r}\dot{\theta} + r^2\ddot{\theta} - r^2 \sin \theta \cos \theta \dot{\phi}^2 = 0$$

Dividing through by r^2 and comparing with 0.2 we get

$$(0.5) \quad \Gamma_{\phi\phi}^{\theta} = -\sin \theta \cos \theta$$

$$(0.6) \quad \Gamma_{r\theta}^{\theta} = \Gamma_{\theta r}^{\theta} = \frac{1}{r}$$

$$(0.7) \quad \Gamma_{\theta\phi}^{\theta} = \Gamma_{\phi\theta}^{\theta} = \Gamma_{\theta\theta}^{\theta} = 0$$

With $a = \phi$ we have

$$(0.8) \quad 2r \sin^2 \theta \dot{r} \dot{\phi} + 2r^2 \sin \theta \cos \theta \dot{\theta} \dot{\phi} + r^2 \sin^2 \theta \ddot{\phi} = 0$$

$$(0.9) \quad \frac{2}{r} \dot{r} \dot{\phi} + 2 \cot \theta \dot{\theta} \dot{\phi} + \ddot{\phi} = 0$$

$$(0.10) \quad \Gamma_{\theta\phi}^{\phi} = \Gamma_{\phi\theta}^{\phi} = \cot \theta$$

$$(0.11) \quad \Gamma_{r\phi}^{\phi} = \Gamma_{\phi r}^{\phi} = \frac{1}{r}$$

$$(0.12) \quad \Gamma_{\theta\theta}^{\phi} = \Gamma_{\phi\phi}^{\phi} = 0$$

For $a = r$, we have

$$(0.13) \quad \ddot{r} - \frac{1}{2} (2r\dot{\theta}^2 + 2r\sin^2 \theta \dot{\phi}^2) = 0$$

$$(0.14) \quad \ddot{r} - r\dot{\theta}^2 - r\sin^2 \theta \dot{\phi}^2 = 0$$

$$(0.15) \quad \Gamma_{\theta\theta}^r = -r$$

$$(0.16) \quad \Gamma_{\phi\phi}^r = -r\sin^2 \theta$$

We can use these results to get the Riemann tensor.

$$(0.17) \quad R_{abcd} = g_{af} R_{bcd}^f$$

$$(0.18) \quad = g_{af} \left(\partial_c \Gamma_{db}^f - \partial_d \Gamma_{cb}^f + \Gamma_{db}^k \Gamma_{ck}^f - \Gamma_{cb}^k \Gamma_{kd}^f \right)$$

Although we know there is only one independent component in 2-d, we can work out all four non-zero components to see how the calculations go.

$$(0.19) \quad R_{\theta\phi\theta\phi} = g_{\theta f} R_{\phi\theta\phi}^f$$

$$(0.20) \quad = g_{\theta\theta} R_{\phi\theta\phi}^{\theta}$$

$$(0.21) \quad = r^2 \left(\partial_{\theta} \Gamma_{\phi\phi}^{\theta} - \partial_{\phi} \Gamma_{\theta\phi}^{\theta} + \Gamma_{\phi\phi}^k \Gamma_{\theta k}^{\theta} - \Gamma_{\theta\phi}^k \Gamma_{k\phi}^{\theta} \right)$$

$$(0.22) \quad = r^2 \left(\sin^2 \theta - \cos^2 \theta - 0 - \frac{r \sin^2 \theta}{r} + \cos^2 \theta \right)$$

$$(0.23) \quad = 0$$

The other components of the Riemann tensor can be evaluated in the same way, and they all come out to zero, so spherical coordinates represent flat 3-d space.