

## GRAVITOELECTRIC AND GRAVITOMAGNETIC DENSITIES FOR THE VACUUM

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 22, Problem 22.2.

The gravitoelectric, gravitomagnetic and curvature densities are

$$(0.1) \quad \rho_g = 2T_{tt} - \eta_{tt}T$$

$$(0.2) \quad \Pi_i = -T_{ti} + \frac{1}{2}\eta_{ti}T$$

$$(0.3) \quad \rho_c = 2T_{ii} - \eta_{ii}T$$

where  $i$  is a spatial index.

The vacuum stress-energy tensor is given in terms of the metric and cosmological constant as

$$(0.4) \quad T_{ij,vac} = -g_{ij} \frac{\Lambda}{8\pi G}$$

Since  $\frac{\Lambda}{8\pi G}$  is very small and  $g_{ij} = \eta_{ij} + h_{ij}$  for a perturbation  $h_{ij}$ , we can approximate:

$$(0.5) \quad T_{ij,vac} \approx -\eta_{ij} \frac{\Lambda}{8\pi G}$$

$$(0.6) \quad T = -\eta_{ij}\eta^{ij} \frac{\Lambda}{8\pi G}$$

$$(0.7) \quad = -\frac{\Lambda}{2\pi G}$$

Therefore

$$(0.8) \quad \rho_g = -\frac{\Lambda}{\pi G} \eta_{tt} \left( -\frac{1}{4} + \frac{1}{2} \right)$$

$$(0.9) \quad = -\frac{\Lambda}{4\pi G}$$

$$(0.10) \quad \Pi_i = \eta_{ti} \frac{\Lambda}{8\pi G} + \eta_{ti} T$$

$$(0.11) \quad = 0$$

$$(0.12) \quad \rho_c = -\frac{\Lambda}{\pi G} \eta_{ii} \left( \frac{1}{4} - \frac{1}{2} \right)$$

$$(0.13) \quad = \frac{\Lambda}{4\pi G}$$

In the third line  $\eta_{ti} = 0$  if  $i$  is a spatial index, and in the fifth line  $\eta_{ii} = +1$ .