

KLEIN-GORDON SOLUTIONS FROM HARMONIC OSCILLATOR

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Michael E. Peskin & Daniel V. Schroeder, *An Introduction to Quantum Field Theory*, (Perseus Books, 1995) - Chapter 2.

Another way of deriving the Klein-Gordon field and its conjugate momentum density is by drawing an analogy with the harmonic oscillator. The idea is to start with the classical Klein-Gordon field $\phi(\mathbf{x}, t)$ and expand it by using a Fourier transform:

$$(1) \quad \phi(\mathbf{x}, t) = \frac{1}{(2\pi)^3} \int d^3 p e^{i\mathbf{p}\cdot\mathbf{x}} \phi(\mathbf{p}, t)$$

Since $\phi(\mathbf{x}, t)$ is a classical field, it must be real, which we can ensure by requiring that $\phi^*(\mathbf{p}, t) = \phi(-\mathbf{p}, t)$. Plugging this into the Klein-Gordon equation gives

$$(2) \quad \partial_\mu \partial^\mu \phi(\mathbf{x}, t) + m^2 \phi(\mathbf{x}, t) = 0$$

$$(3) \quad \frac{1}{(2\pi)^3} \int d^3 p \left[\frac{\partial^2}{\partial t^2} - \nabla^2 + m^2 \right] e^{i\mathbf{p}\cdot\mathbf{x}} \phi(\mathbf{p}, t) = 0$$

$$(4) \quad \frac{1}{(2\pi)^3} \int d^3 p e^{i\mathbf{p}\cdot\mathbf{x}} \left[\frac{\partial^2}{\partial t^2} + |\mathbf{p}|^2 + m^2 \right] \phi(\mathbf{p}, t) = 0$$

Since this must be true for all $\phi(\mathbf{p}, t)$, the integrand must be zero so we get

$$(5) \quad \left[\frac{\partial^2}{\partial t^2} + |\mathbf{p}|^2 + m^2 \right] \phi(\mathbf{p}, t) = 0$$

The classical equation of motion for a harmonic oscillator in one dimension is

$$(6) \quad F = -kx$$

$$(7) \quad m_0 \frac{d^2 x}{dt^2} + kx = 0$$

The solution is an oscillator with frequency

$$(8) \quad \omega = \sqrt{\frac{k}{m_0}}$$

The corresponding Hamiltonian is

$$(9) \quad H = \frac{p^2}{2m_0} + \frac{1}{2}kx^2$$

Comparing this with 5, we see that the Klein-Gordon equation in momentum space has the same form as a harmonic oscillator with x replaced by \mathbf{p} , $k = |\mathbf{p}|^2 + m^2$ and $m_0 = 1$, so its solution is an oscillator with frequency

$$(10) \quad \omega_{\mathbf{p}} = \sqrt{|\mathbf{p}|^2 + m^2}$$

In the algebraic solution of the harmonic oscillator in non-relativistic quantum mechanics, we introduced the raising and lowering operators (renaming $a_+ = a^\dagger$ and $a_- = a$ to be consistent with P&S's notation):

$$(11) \quad a^\dagger = \frac{1}{\sqrt{2\hbar m_0 \omega}} [-ip + m_0 \omega x]$$

$$(12) \quad a = \frac{1}{\sqrt{2\hbar m_0 \omega}} [ip + m_0 \omega x]$$

In natural units $\hbar = 1$ and with $m_0 = 1$ this gives

$$(13) \quad a^\dagger = \frac{1}{\sqrt{2\omega}} [-ip + \omega x]$$

$$(14) \quad a = \frac{1}{\sqrt{2\omega}} [ip + \omega x]$$

which can be inverted to give

$$(15) \quad x = \frac{1}{\sqrt{2\omega}} (a + a^\dagger)$$

$$(16) \quad p = -i\sqrt{\frac{\omega}{2}} (a - a^\dagger)$$

Using the commutation relation

$$(17) \quad [x, p] = i$$

we get

$$(18) \quad [a, a^\dagger] = 1$$

and the Hamiltonian comes out to

$$(19) \quad H_{SHO} = \omega \left(a^\dagger a + \frac{1}{2} \right)$$

The effects of a and a^\dagger are to lower and raise a state by an energy quantum ω . The wave functions are

$$(20) \quad a |n\rangle = \sqrt{n} |n-1\rangle$$

$$(21) \quad a^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle$$

To apply all this to the quantum Klein-Gordon equation, we want to interpret x in 15 as a quantum field $\phi(\mathbf{p})$, and leave p in 16 as it is. The field $\phi(\mathbf{x})$ in position space is the integral 1 over $\phi(\mathbf{p})$ in momentum space, with each momentum \mathbf{p} contributing its own raising and lowering operators $a_{\mathbf{p}}$ and $a_{-\mathbf{p}}^\dagger$ from 13 and 14. That is, we get

$$(22) \quad \phi(\mathbf{x}) = \frac{1}{(2\pi)^3} \int d^3 p e^{i\mathbf{p}\cdot\mathbf{x}} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (a_{\mathbf{p}} + a_{-\mathbf{p}}^\dagger)$$

$$(23) \quad \pi(\mathbf{x}) = \frac{-i}{(2\pi)^3} \int d^3 p e^{i\mathbf{p}\cdot\mathbf{x}} \sqrt{\frac{\omega_{\mathbf{p}}}{2}} (a_{\mathbf{p}} - a_{-\mathbf{p}}^\dagger)$$

We can invert these equations to get expressions for a and a^\dagger . We multiply both sides by $e^{-i\mathbf{p}'\cdot\mathbf{x}}$ and integrate over $d^3 x$:

$$(24) \quad \int d^3 x e^{-i\mathbf{p}'\cdot\mathbf{x}} \phi(\mathbf{x}) = \int d^3 p \left(\frac{1}{(2\pi)^3} \int d^3 x e^{i(\mathbf{p}-\mathbf{p}')\cdot\mathbf{x}} \right) \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (a_{\mathbf{p}} + a_{-\mathbf{p}}^\dagger)$$

$$(25) \quad = \int d^3 p \delta(\mathbf{p} - \mathbf{p}') \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (a_{\mathbf{p}} + a_{-\mathbf{p}}^\dagger)$$

$$(26) \quad = \frac{1}{\sqrt{2\omega_{\mathbf{p}'}}} (a_{\mathbf{p}'} + a_{-\mathbf{p}'}^\dagger)$$

$$(27) \quad \int d^3 x e^{-i\mathbf{p}'\cdot\mathbf{x}} \pi(\mathbf{x}) = -i \sqrt{\frac{\omega_{\mathbf{p}'}}{2}} (a_{\mathbf{p}'} - a_{-\mathbf{p}'}^\dagger)$$

Dropping the primes on \mathbf{p}' and solving for a and a^\dagger we get (since $\omega_{\mathbf{p}} = \omega_{-\mathbf{p}}$):

$$(28) \quad a_{\mathbf{p}} = \int d^3x \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} e^{-i\mathbf{p}\cdot\mathbf{x}} [i\pi(\mathbf{x}) + \omega_{\mathbf{p}}\phi(\mathbf{x})]$$

$$(29) \quad a_{\mathbf{p}}^\dagger = \int d^3x \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} e^{i\mathbf{p}\cdot\mathbf{x}} [-i\pi(\mathbf{x}) + \omega_{\mathbf{p}}\phi(\mathbf{x})]$$

PINGBACKS

Pingback: Klein-Gordon equation from harmonic oscillator: Hamiltonian, creation and

Pingback: Klein-Gordon field in position space

Pingback: Klein-Gordon field in the Heisenberg picture; time dependence