

HARMONIC OSCILLATOR - MATRIX ELEMENTS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 29 July 2021.

In analyzing the harmonic oscillator, we used the raising and lowering operators to calculate $\langle x \rangle$ and $\langle p \rangle$, finding that they are both zero for all stationary states. These quantities are really the diagonal elements of the matrices X and P . That is

$$\langle x \rangle_{nn} = \langle n|x|n \rangle \quad (1)$$

$$= X_{nn} \quad (2)$$

We can use the same technique to calculate the off-diagonal elements.

We review the equations involving the raising and lowering operators first:

$$x = \sqrt{\frac{\hbar}{2m\omega}}(a_+ + a_-) \quad (3)$$

$$p = i\sqrt{\frac{\hbar m\omega}{2}}(a_+ - a_-) \quad (4)$$

$$a_+\psi_n = \sqrt{n+1}\psi_{n+1} \quad (5)$$

$$a_-\psi_n = \sqrt{n}\psi_{n-1} \quad (6)$$

The general matrix elements for the operator x can then be calculated:

$$\langle n|x|n' \rangle = \sqrt{\frac{\hbar}{2m\omega}}(\sqrt{n'+1}\langle n|n'+1 \rangle + \sqrt{n'}\langle n|n'-1 \rangle) \quad (7)$$

$$= \sqrt{\frac{\hbar}{2m\omega}}(\sqrt{n'+1}\delta_{n,n'+1} + \sqrt{n'}\delta_{n,n'-1}) \quad (8)$$

By similar reasoning we get the matrix elements for p :

$$\langle n|p|n' \rangle = i\sqrt{\frac{\hbar m\omega}{2}}(\sqrt{n'+1}\delta_{n,n'+1} - \sqrt{n'}\delta_{n,n'-1}) \quad (9)$$

These results agree with those found by doing the integrals involving Hermite polynomials.

We now have all the matrix elements of X and P so it would be interesting to calculate the full hamiltonian matrix, which is

$$H = \frac{1}{2m}P^2 + \frac{m\omega^2}{2}X^2 \quad (10)$$

In order to calculate the squares of the two matrices, we observe that both X and P are *tridiagonal* matrices with the added condition that their main diagonals are all zero. That is, the two diagonals above and below the main diagonal are the only places with non-zero elements. The square of such a matrix will have non-zero elements only on the main diagonal, and on the diagonals *two* above and below the main diagonal (you can verify this by drawing out such a matrix and seeing where the non-zero elements lie, or by doing tedious calculations with indices).

We can demonstrate how these elements can be calculated by considering the diagonal elements of X^2 .

$$X_{nn}^2 = \sum_{n'} \langle n|x|n'\rangle \langle n'|x|n\rangle \quad (11)$$

$$= \frac{\hbar}{2m\omega} \sum_{n'} [\sqrt{n'+1}\delta_{n,n'+1} + \sqrt{n'}\delta_{n,n'-1}] [\sqrt{n+1}\delta_{n',n+1} + \sqrt{n}\delta_{n',n-1}] \quad (12)$$

$$= \frac{\hbar}{2m\omega} (2n+1) \quad (13)$$

The last line is obtained by noting that all the terms in the sum contain the product of two Kronecker deltas, so only in those cases where *both* deltas are non-zero is there a non-zero contribution to the sum. This happens only in the terms involving the product of the first and fourth terms (where $n' = n - 1$) and the second and third terms (where $n' = n + 1$).

By a similar argument, we get

$$P_{nn}^2 = \frac{\hbar m\omega}{2} (2n+1) \quad (14)$$

Therefore the *diagonal* elements of $(1/2m)P^2 + (m\omega^2/2)X^2$ are

$$H_{nn} = \hbar\omega \left(n + \frac{1}{2} \right) \quad (15)$$

which is what you would expect, as these are the energy levels of the harmonic oscillator.

It remains only to show that the off-diagonal elements of H are zero.

$$X_{nm}^2 = \sum_{n'} \langle n|x|n' \rangle \langle n'|x|m \rangle \quad (16)$$

$$= \frac{\hbar}{2m\omega} \sum_{n'} [\sqrt{n'+1} \delta_{n,n'+1} + \sqrt{n'} \delta_{n,n'-1}] [\sqrt{m+1} \delta_{n',m+1} + \sqrt{m} \delta_{n',m-1}] \quad (17)$$

To see which non-zero elements exist on row n , we note that for a given value of n , we must have either $n' = n - 1$ or $n' = n + 1$ in order for one of the deltas in the first term to be non-zero. If $n' = n - 1$, then in the second term, we must have either $n - 1 = m + 1$ or $n - 1 = m - 1$. The second case results in a diagonal element which we have already considered, so we need consider only the case $m = n - 2$. In this case, the matrix element is

$$X_{n,n-2}^2 = \frac{\hbar}{2m\omega} \sqrt{n(n-1)} \quad (18)$$

Similarly, if $n' = n + 1$, the non-diagonal term is $n + 1 = m - 1$ or $m = n + 2$, and we get

$$X_{n,n+2}^2 = \frac{\hbar}{2m\omega} \sqrt{(n+1)(n+2)} \quad (19)$$

Similar reasoning gives us the elements from P^2 :

$$P_{n,n-2}^2 = -\frac{\hbar m\omega}{2} \sqrt{n(n-1)} \quad (20)$$

$$P_{n,n+2}^2 = -\frac{\hbar m\omega}{2} \sqrt{(n+1)(n+2)} \quad (21)$$

Combining these two results, we see that the *non-diagonal* elements of $(1/2m)P^2 + (m\omega^2/2)X^2$ are all zero.

PINGBACKS

Pingback: Second order non-degenerate perturbation theory