

HARMONIC OSCILLATOR EXCITED STATES - NUMERICAL SOLUTION

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 15 June 2021.

In the last post, we looked at calculating the ground state energy of the harmonic oscillator numerically, using the 'wag the dog' method. In this post, we'll look at calculating the first three excited states in the same way.

In the analytic solution, we arrived at the following differential equation

$$\frac{d^2\psi}{dy^2} = (y^2 - \epsilon)\psi \quad (1)$$

where we've used dimensionless parameters

$$y \equiv \sqrt{\frac{m\omega}{\hbar}}x \quad (2)$$

$$\epsilon \equiv \frac{2E}{\hbar\omega} \quad (3)$$

The ground state occurs when $\epsilon = 1$ and the next three excited states occur when $\epsilon = 3, 5, 7$ respectively. We know this, of course, only because we've solved the problem analytically, but for the purposes of numerical solution, we will show what happens if we take ϵ slightly less and greater than the correct values in each case. We'll use the same Maple code as in the previous post.

The solutions for $\epsilon = 3, 7$ are odd, so as we saw in the last post, we'll need to take $\psi(0) = 0$ and $\psi'(0) \neq 0$. Just for fun, we'll take $\psi'(0) = -1$. With $\epsilon = 2.999$, we get Fig. 1.

For $\epsilon = 3.001$ we get Fig. 2.

The two tails of the function have flipped, but the central region remains the same since it is constrained by the initial conditions. Finally, with $\epsilon = 3.000$ we get the correct solution, as in Fig. 3.




FIGURE 1.




FIGURE 2.

For the second excited state, the solution is even so we take $\psi(0) = -1$ (again, just for variety - the actual numerical value doesn't matter since any multiple of a solution is also a solution, and the *actual* value of $\psi(0)$ would




FIGURE 3.

be determined by normalization anyway) and $\psi'(0) = 0$. We try $\epsilon = 4.999$ and get Fig. 4.

For $\epsilon = 5.001$ we have Fig. 5.

And finally for $\epsilon = 5.000$ we get the correct solution, as in Fig. 6.

For the third excited state, the solution is again odd, so we revert to $\psi(0) = 0$ and $\psi'(0) = -1$. With $\epsilon = 6.999$ we have Fig. 7.

For $\epsilon = 7.001$ we have Fig. 8.

And finally for $\epsilon = 7.000$ we get the correct solution in Fig. 9.

PINGBACKS

Pingback: Infinite square well - numerical solution

Pingback: Finite square well - numerical solution




FIGURE 4.




FIGURE 5.




FIGURE 6.




FIGURE 7.




FIGURE 8.




FIGURE 9.