

SPIN MATRICES - GENERAL CASE

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 13 September 2021.

We can generalize the calculation for spin 3/2 to get the spin matrices for any spin s .

We first find S_{\pm} from the equation

$$S_{\pm}|sm\rangle = \hbar\sqrt{s(s+1) - m(m\pm 1)}|sm\pm 1\rangle \quad (1)$$

First, consider S_+ . We know that $S_+|ss\rangle = 0$ and that $m = s - k$ for $k = 1 \dots 2s$ for the remaining eigenstates of S_z . We can also see from the spin 3/2 case that the S_+ matrix contains non-zero entries only on the diagonal above the main diagonal (since the raising operator always raises an eigenstate m of S_z to the state $m + 1$), and that these entries consist of the square root factor given in 1 for values of m starting with $m = s - 1$ as matrix element $S_{+(1,2)}$ (i.e. the element in row 1, column 2 of matrix S_+), and then working down the diagonal, with element $S_{+(k,k+1)}$ being the square root factor for $m = s - k$.

Letting $m = s - k$ for $k = 1 \dots 2s$ we get

$$\sqrt{s(s+1) - m(m+1)} = \sqrt{s(s+1) - (s-k)(s-k+1)} \quad (2)$$

$$= \sqrt{2sk + k(1-k)} \quad (3)$$

$$\equiv c_k \quad (4)$$

We can therefore write

$$S_+ = \hbar \begin{bmatrix} 0 & c_1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & c_2 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & c_3 & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 0 & c_{2s} \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix} \quad (5)$$

For those readers wanting to get the answer as given in Griffiths's book, we can make the substitution

$$j \equiv -k + 1 + s \quad (6)$$

so that $j = s \dots -s + 1$ as $k = 1 \dots 2s$. Then

$$\sqrt{2sk + k(1-k)} = \sqrt{2(1+s-j)s + (1+s-j)(j-s)} \quad (7)$$

$$= \sqrt{(1+s-j)(j+s)} \quad (8)$$

$$\equiv b_j \quad (9)$$

as given in the book.

The calculation for S_- is similar, giving $S_{-(k+1,k)} = c_k$ for $k = 1 \dots 2s$, so we get

$$S_- = \hbar \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ c_1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & c_2 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & c_3 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & c_{2s} & 0 \end{bmatrix} \quad (10)$$

In Griffiths's notation, $S_{-(k+1,k)} = b_j$, where $j = -k + 1 + s$ as before, and we can then use the formulas $S_x = (S_+ + S_-)/2$ and $S_y = (S_+ - S_-)/2i$ to get the actual spin matrices, which will have non-zero entries on the two diagonals on either side of the main diagonal.