

## COVARIANT DERIVATIVE OF THE METRIC TENSOR

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Post date: 20 Jan 2023.

One interesting and useful theorem is that the covariant derivative of *any* metric tensor is always zero. We can show this by using the expression for the covariant derivative of a general tensor to say:

$$\nabla_j g_{kl} = \partial_j g_{kl} - \Gamma_{jk}^m g_{ml} - \Gamma_{jl}^m g_{km} \quad (1)$$

We can combine this with the explicit expression for the Christoffel symbols:

$$\Gamma_{ij}^m = \frac{1}{2} g^{ml} (\partial_j g_{il} + \partial_i g_{lj} - \partial_l g_{ji}) \quad (2)$$

Substituting, we get

$$\nabla_j g_{kl} = \partial_j g_{kl} - \frac{1}{2} g^{mn} (\partial_k g_{jn} + \partial_j g_{nk} - \partial_n g_{kj}) g_{ml} - \frac{1}{2} g^{mn} (\partial_l g_{jn} + \partial_j g_{nl} - \partial_n g_{lj}) g_{km} \quad (3)$$

Since  $g^{mn} g_{ml} = \delta_l^n$ , we get

$$\nabla_j g_{kl} = \partial_j g_{kl} - \frac{1}{2} (\partial_k g_{jl} + \partial_j g_{lk} - \partial_l g_{kj}) - \frac{1}{2} (\partial_l g_{jk} + \partial_j g_{kl} - \partial_k g_{lj}) \quad (4)$$

Now we use the symmetry of the metric tensor:  $g_{kl} = g_{lk}$ :

$$\nabla_j g_{kl} = \partial_j g_{lk} - \frac{1}{2} (\partial_k g_{jl} + \partial_j g_{lk} - \partial_l g_{kj}) - \frac{1}{2} (\partial_l g_{kj} + \partial_j g_{lk} - \partial_k g_{jl}) \quad (5)$$

$$= 0 \quad (6)$$

### PINGBACKS

Pingback: Covariant derivative of the metric tensor - application to a coordinate

Pingback: Metric tensor as a stress-energy tensor

Pingback: Conservation of four-momentum implies the geodesic equation

Pingback: Einstein equation solution for the interior of a spherically symmetric star