

ELECTROMAGNETIC FIELD TENSOR - A COUPLE OF MAXWELL'S EQUATIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 25 Aug 2022.

The electromagnetic field tensor F^{ij} is

$$F^{ij} = \begin{bmatrix} 0 & E_x & E_y & E_z \\ -E_x & 0 & B_z & -B_y \\ -E_y & -B_z & 0 & B_x \\ -E_z & B_y & -B_x & 0 \end{bmatrix} \quad (1)$$

The various electromagnetic laws can be expressed by the equation

$$\partial_i F_{jk} + \partial_k F_{ij} + \partial_j F_{ki} = 0 \quad (2)$$

The lowered version F_{ij} of F^{ij} in the flat metric of special relativity is

$$F_{ij} = \begin{bmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & B_z & -B_y \\ E_y & -B_z & 0 & B_x \\ E_z & B_y & -B_x & 0 \end{bmatrix} \quad (3)$$

Because $F_{ij} = -F_{ji}$, if we set two of the indices equal in 2, the LHS is identically zero. For example if $i = j$:

$$\partial_i F_{ik} + \partial_k F_{ii} + \partial_i F_{ki} = -\partial_i F_{ki} + 0 + \partial_i F_{ki} \quad (4)$$

$$= 0 \quad (5)$$

Since the three terms are cyclic permutations of each other, setting any other pair of indices equal gives the same result.

If we choose $i = y, j = x, k = z$, we get

$$-\partial_y B_y - \partial_z B_z - \partial_x B_x = 0 \quad (6)$$

This is the law $\nabla \cdot \mathbf{B} = 0$.

If we now choose $i = y, j = x, k = t$, we get

$$-\partial_y E_x + \partial_t B_z + \partial_x E_y = 0 \quad (7)$$

This is the z component of Faraday's law $\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$. Other choices give the remaining components of Faraday's law.

PINGBACKS

Pingback: [Electromagnetic field tensor - cyclic derivative relation](#)