

METRIC TENSOR IN PARABOLIC COORDINATES

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 23 June 2021.

Another example of a non-rectangular 2-d curved coordinate system. This time we have a parabolic bowl with equation

$$z = br^2 \quad (1)$$

where $r^2 = x^2 + y^2$ and b is a constant. We use the two coordinates r (as defined here) and the azimuthal angle ϕ .

Using the same technique as with the sphere, we consider infinitesimal displacements along the constant curves. The curve of constant ϕ is a parabola, while the curve of constant r is a circle at height $z = br^2$. The tangents to the two curves at a given point are always perpendicular, so the metric g_{ij} will be diagonal. To find the diagonal components, consider an infinitesimal displacement ds . We have

$$ds = dr\mathbf{e}_r + d\phi\mathbf{e}_\phi \quad (2)$$

and our job is to find the lengths of the two basis vectors.

Consider first a displacement along \mathbf{e}_r . As r increases, we move a distance up the side of the parabolic bowl. This displacement consists of a horizontal increment (that is, radially outward from the z axis) of size dr and a vertical increment of size $dz = 2br dr$. By Pythagoras, the total displacement is $ds = \sqrt{1 + (2br)^2} dr$. The length of \mathbf{e}_r is therefore $\sqrt{1 + (2br)^2}$.

A displacement along \mathbf{e}_ϕ is a displacement around a circle of constant r , so here $ds = r d\phi$, giving the length of \mathbf{e}_ϕ as r . The metric is then

$$g_{ij} = \begin{bmatrix} 1 + (2br)^2 & 0 \\ 0 & r^2 \end{bmatrix} \quad (3)$$

PINGBACKS

Pingback: Geodesic equation on a paraboloid