

SCHWARZSCHILD METRIC - GRAVITATIONAL RED SHIFT

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Post date: 29 Aug 2022.

The relation between the proper time interval and Schwarzschild time interval for an object at rest is

$$\Delta\tau = \left(1 - \frac{2GM}{r}\right)^{1/2} \Delta t \quad (1)$$

We can use this relation to derive a formula for the gravitational red shift. The key to this is that the wavelength of light is $\lambda = c\Delta\tau$, where $c = 1$ is the speed of light and $\Delta\tau$ is the time interval as measured by an observer required for a single wavelength to be emitted or received. Why $\Delta\tau$ instead of Δt ? As far as I understand it, this is because τ is the only correct measure of time for an object at rest. The Schwarzschild t coordinate, although it's called the 'time coordinate', isn't really a measure of time directly.

The red shift arises because if we emit a light beam at $r = r_E$ in a direction radially outwards from the mass M and receive the light beam at $r = r_R > r_E$, then the interval Δt required for the passage of a single wavelength must be the same at both the emitter and the receiver. Why? Because the metric doesn't depend on t .

The proper time interval, however, is not the same at the two points because of the relation above. Plugging in the values, we get

$$\frac{\lambda_R}{\lambda_E} = \frac{\Delta\tau_R}{\Delta\tau_E} \quad (2)$$

$$= \sqrt{\frac{1 - 2GM/r_R}{1 - 2GM/r_E}} \quad (3)$$

This is the gravitational red shift formula. For $r_R \rightarrow \infty$, the formula reduces to

$$\frac{\lambda_R}{\lambda_E} = \frac{1}{\sqrt{1 - 2GM/r_E}} \quad (4)$$

This formula could be used, for example, to calculate the red shift due to a star when observed from Earth.

If both distances are large compared to $2GM$, we can expand the formula in a series up to first order:

$$\frac{\lambda_R}{\lambda_E} \approx \left(1 - \frac{GM}{r_R}\right) \left(1 + \frac{GM}{r_E}\right) \quad (5)$$

$$= 1 + GM \left(\frac{1}{r_E} - \frac{1}{r_R}\right) + \dots \quad (6)$$

As a further approximation, if the distance $h = r_R - r_E \ll r_E$, that is, the distance between emission and reception is small compared with the radial coordinate, then we can write

$$\frac{\lambda_R}{\lambda_E} \approx 1 + GM \left(\frac{1}{r_E} - \frac{1}{r_R}\right) \quad (7)$$

$$= 1 + GM \left(\frac{h}{r_E r_R}\right) \quad (8)$$

$$\approx 1 + \frac{GM}{r^2} h \quad (9)$$

where r in the last line can be taken as the average of r_R and r_E . In this limit, we'd expect Newton's law of gravitation to apply, and a particle a distance r from a mass M experiences an acceleration $g = GM/r^2$, so we have

$$\frac{\lambda_R}{\lambda_E} \approx 1 + gh \quad (10)$$

Example 1. Suppose we have a neutron star with mass $M = 3 \times 10^{30}$ kg and Schwarzschild radial coordinate at the surface of $r_E = 1.2 \times 10^4$ m. The red shift observed by a satellite orbiting the star at a radius $r_R = 1.7 \times 10^4$ m can be calculated using the approximation formula. We need to express G in relativistic units (that is, where $c = 1$ so that GM has the units of length). Since the units of G are $\text{m}^3 \text{kg}^{-1} \text{s}^{-2}$, we need to eliminate the reference to seconds which we can do by dividing by $c^2 = 9 \times 10^{16} \text{m}^2 \text{s}^{-2}$. That is

$$G = \frac{6.67 \times 10^{-11}}{9 \times 10^{16}} \quad (11)$$

$$= 7.41 \times 10^{-28} \text{m kg}^{-1} \quad (12)$$

For the neutron star,

$$GM = (7.41 \times 10^{-28}) (3 \times 10^{30}) \quad (13)$$

$$= 2.223 \times 10^3 \text{ m} \quad (14)$$

We take

$$r = \frac{1}{2} (r_E + r_R) \quad (15)$$

$$= 1.45 \times 10^4 \text{ m} \quad (16)$$

$$g = \frac{GM}{r^2} \quad (17)$$

$$= \frac{2.223 \times 10^3}{(1.45 \times 10^4)^2} \quad (18)$$

$$= 1.06 \times 10^{-5} \text{ m}^{-1} \quad (19)$$

Incidentally, this is a massive acceleration compared to that on the Earth's surface. In SI units, this comes out to $(1.06 \times 10^{-5}) (9 \times 10^{16}) = 9.54 \times 10^{11} \text{ m s}^{-2}$.

The fractional red shift is

$$\frac{\lambda_R - \lambda_E}{\lambda_E} \approx g(r_R - r_E) \quad (20)$$

$$= (1.06 \times 10^{-5}) (5 \times 10^3) \quad (21)$$

$$= 0.0529 \quad (22)$$

The exact value is

$$\frac{\lambda_R - \lambda_E}{\lambda_E} = \sqrt{\frac{1 - 2GM/r_R}{1 - 2GM/r_E}} - 1 \quad (23)$$

$$= 0.0831 \quad (24)$$

PINGBACKS

Pingback: Schwarzschild metric - redshift of Sirius B

Pingback: Red-shifts and blue-shifts

Pingback: Gravitational redshift from the Killing vector