

STRESS-ENERGY TENSOR FOR A PERFECT FLUID AT REST

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 3 Feb 2021.

We've seen the stress-energy tensor T^{ij} for the special case of 'dust', that is a system in which we have a collection of particles that are all at rest relative to each other. In that special case,

$$T^{ij} \equiv \rho_0 u^i u^j \quad (1)$$

where ρ_0 is the energy density of the dust in its rest frame, and u^i is the four-velocity of a local inertial frame (LIF) moving relative to the dust.

Real fluids, of course, contain particles that are in some sort of random motion, even if the fluid as a whole is at rest. We can derive the stress-energy tensor for such fluids if we assume that they are *perfect* or *ideal* fluids, which means that the particles within the fluid don't interact with each other. This is still a simplification since in all real fluids, even gases, the particles do interact. In liquids, this interaction gives the fluids viscosity.

The idea is that we can build up a description of a perfect fluid by considering subsets of the particles where within each subset, the particles act like dust, that is, they all move at the same velocity. We can work out the contribution to T^{ij} from each of these subsets and then add them up to get the overall tensor.

To get started, consider the subset of particles that have four-velocity u^i as measured in the overall rest frame of the fluid (that is, the frame in which the average of the velocities of all the particles is zero). Then the contribution to T^{ij} from that subset is

$$\Delta T^{ij} = \rho_0 u^i u^j \quad (2)$$

where ρ_0 is now the density of particles within that subset only, and measured in that subset's rest frame. In terms of the number density n_0 (again measured in the subset's frame) this is

$$\Delta T^{ij} = n_0 m u^i u^j \quad (3)$$

Due to length contraction, the number density as measured in the fluid's rest frame is

$$n = \frac{n_0}{\sqrt{1-\beta^2}} = \gamma n_0 = u^t n_0 \quad (4)$$

where u^t is the time component of the four-velocity. Therefore we can write

$$\Delta T^{ij} = \frac{n}{u^t} m u^i u^j = \frac{n m u^i m u^j}{m u^t} = n \frac{p^i p^j}{p^t} \quad (5)$$

where p^i is the four-momentum of a single particle as measured in the fluid's frame.

Now suppose we look at a cubic volume at rest in the fluid's frame with side length L (as measured in the fluid's frame; because of length contraction, the volume won't be a cube in any of the individual particle's frames). Then if N is the total number of particles in the subset, we have $n = N/L^3$ so

$$\Delta T^{ij} = \frac{N}{L^3} \frac{p^i p^j}{p^t} \quad (6)$$

The total stress-energy tensor is then the integral of this quantity over all of momentum space, since we need to add up contributions of all the subsets with each possible momentum. Since there will be different numbers of particles in each subset, the quantity N becomes a function of momentum: $N = N(\mathbf{p})$. If we assume that the motions of the particles are truly random, then it is equally likely that a particle with magnitude of momentum p is moving in any direction, so we'd expect $N(\mathbf{p})$ to be a function of the *magnitude* of \mathbf{p} only, rather than on its direction as well. That is, we have

$$N = N(p) \quad (7)$$

$$p = \sqrt{(p^x)^2 + (p^y)^2 + (p^z)^2} \quad (8)$$

and the total stress-energy tensor is

$$T^{ij} = \frac{1}{L^3} \int \int \int N(p) \frac{p^i p^j}{p^t} dp^x dp^y dp^z \quad (9)$$

Looking at the terms in the integrand, we see that $N(p)$ is an even function of each of p^x , p^y and p^z . Since $p^t = m u^t = \frac{m^2}{m\sqrt{1-\beta^2}} = \frac{m^2}{\sqrt{m^2-p^2}}$, it too is an even function of each of p^x , p^y and p^z . If we consider an off-diagonal element of T^{ij} such as T^{yz} , we have

$$T^{yz} = \frac{1}{L^3} \int dp^x \int p^y dp^y \int p^z \frac{N(p)}{p^t} dp^z \quad (10)$$

The first integral over p^z gives zero, since we are integrating the product of an even function $N(p)/p^t$ and an odd function p^z over the range $-\infty$ to ∞ . (We are also assuming, of course, that $N(p) \rightarrow 0$ as $p \rightarrow \infty$ which is reasonable, since we'd expect the number of particles with very large momenta to go to zero.)

The same argument applies to any off-diagonal element since any such element will integrate an odd function p^i multiplied by the even function $N(p)/p^t$. This argument applies also to off-diagonal elements where one of the indices is t , since the other index will be a spatial index and provide the odd function that multiplies $N(p)/p^t$.

If we look at the diagonal elements, however, we get the product of two even functions, as with T^{zz} :

$$T^{zz} = \frac{1}{L^3} \int dp^x \int dp^y \int (p^z)^2 \frac{N(p)}{p^t} dp^z \quad (11)$$

and this will not, in general, be zero (unless no particles have a z component of momentum). Therefore, for the special case of a perfect fluid at rest, the stress-energy tensor is diagonal.

To interpret these elements we can use a clever little argument as follows. Look at a single particle that bounces elastically (that is, it doesn't lose any energy in a collision) off the walls of the cube. As seen in the fluid's frame, the time taken between bounces off, say, one of the faces perpendicular to the x axis is the total distance in the x direction it travels (which is twice the width of the cube, or $2L$) divided by its x speed:

$$\Delta t_x = \frac{2L}{v^x} \quad (12)$$

Since the collision is elastic, the particle rebounds from the wall with the same magnitude of v_x but the opposite direction, so its x momentum has been exactly reversed. This means that the wall imparted a change in x momentum of

$$\Delta p^x = 2mv^x \quad (13)$$

The pressure is the average force per unit area, so for this single particle, the average force is the change in momentum per unit time, or

$$\Delta F^x = \frac{\Delta p^x}{\Delta t_x} = \frac{mv^x v^x}{L} = \frac{p^x v^x}{L} \quad (14)$$

The pressure is the force per unit area, so

$$\Delta P = \frac{\Delta F^x}{L^2} = \frac{p^x v^x}{L^3} = \frac{p^x \gamma m v^x}{\gamma m} = \frac{(p^x)^2}{p^t L^3} \quad (15)$$

(Note that v^x is the x component of the particle's velocity as measured in the fluid's frame, so the x component of its *four*-velocity is $\gamma v^x = u^x$.)

The total pressure is therefore the integral of this over all particles, which is

$$P = \int \int \int N(p) \frac{(p^x)^2}{p^t L^3} dp^x dp^y dp^z = T^{xx} \quad (16)$$

by comparison with 9.

There's nothing special about choosing the x direction in this calculation, so we would end up with the same result having chosen y or z . Thus

$$T^{xx} = T^{yy} = T^{zz} = P \quad (17)$$

For the remaining component T^{tt} we have

$$T^{tt} = \int \int \int N(p) \frac{(p^t)^2}{p^t L^3} dp^x dp^y dp^z \quad (18)$$

$$= \frac{1}{L^3} \int \int \int N(p) p^t dp^x dp^y dp^z \quad (19)$$

Since p^t is the energy of a particle with four-momentum p^i , this integral represents the total energy density ρ of the fluid. We therefore get, for a perfect fluid in its rest frame:

$$T^{ij} = \begin{bmatrix} \rho & 0 & 0 & 0 \\ 0 & P & 0 & 0 \\ 0 & 0 & P & 0 \\ 0 & 0 & 0 & P \end{bmatrix} \quad (20)$$

Now suppose we want to generalize this to the case where the fluid is viewed from some more general coordinate system, possibly in curved spacetime. The result is simply stated in every source I could find. The result is

$$T^{ij} = (\rho_0 + P_0) u^i u^j + P_0 g^{ij} \quad (21)$$

where ρ_0 is the energy density of the fluid and P_0 is the pressure, both measured in the fluid's rest frame. We can verify that this reduces to the form 20 when viewed in the fluid's rest frame, since in that frame, $g^{ij} = \eta^{ij} = [-1, 1, 1, 1]$ and the four-velocity is $u^i = [1, 0, 0, 0]$, so all off-diagonal elements are automatically zero, and

$$T^{tt}(\text{rest frame}) = (\rho_0 + P_0)u^t u^t - P_0 = \rho_0 + P_0 - P_0 = \rho_0 \quad (22)$$

$$T^{xx}(\text{rest frame}) = T^{yy} = T^{zz} = (\rho_0 + P_0) \times 0 + P_0 \times 1 = P_0 \quad (23)$$

$$T^{ij} = \begin{bmatrix} \rho_0 & 0 & 0 & 0 \\ 0 & P_0 & 0 & 0 \\ 0 & 0 & P_0 & 0 \\ 0 & 0 & 0 & P_0 \end{bmatrix} \quad (24)$$

The justification of 21 appears to be that it is a proper tensor equation (since u^i is a four-vector and g^{ij} is a tensor, and ρ_0 and P_0 are both scalars), so it is valid in all coordinate systems. The fact that it reduces to the earlier form in the fluid's rest frame shows that it has the right form in that coordinate system. What isn't clear to me is that this is the *only* tensor that reduces to the correct form. Anyone who knows of a proper derivation, please leave a comment.

COMMENTS

11 Oct 2023, 02.23.

Hi, this is John B. Wilson of SciRealm.org

In your derivation of the Perfect Fluid Stress-Energy Tensor, you start with the Dust form with (1) independent parameter: energy density and build up to the Perfect Fluid form with (2) independent parameters: energy density & pressure.

Perhaps a better way would be to consider the top-down approach.

A general symmetric 4D 2-index tensor has 10 independent parameters.

These (10) parameters make up the components of a Relativistic Fluid, which can have heat (3) and anisotropic stress (5) properties in addition to the energy density (1) and isotropic pressure (1).

Then, taking special cases, setting heat and stress properties to zero, one can go down to the perfect fluid case.

Then, taking a further special case, setting the pressure to zero, one can go down to the dust case.

Alternately, one could set the pressure to other certain values to get the PhotonGas=Radiation Fluid or LambdaVacuum cases.

See: <http://scirealm.org/SRQM.pdf> pg. 44 at time of post

PINGBACKS

Pingback: Stress-energy tensor - conservation equations

Pingback: Stress-energy tensor - symmetry

Pingback: Stress-energy tensor for perfect fluid - general coordinates

Pingback: Stress-energy tensor at the centre of the sun

Pingback: [Stress-energy tensor - relativistic perfect fluid](#)

Pingback: [Stress-energy tensor for a photon gas](#)

Pingback: [Metric tensor as a stress-energy tensor](#)

Pingback: [Stress-energy tensor in a local orthonormal frame](#)

Pingback: [Stress-energy tensor - negative pressure revisited](#)

Pingback: [Dominant energy condition](#)

Pingback: [Einstein equation in the Newtonian limit](#)

Pingback: [Einstein equation for an exponential metric](#)