

## HEAT CAPACITY OF A SINGLE WATER MOLECULE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Daniel V. Schroeder, *An Introduction to Thermal Physics*, (Addison-Wesley, 2000) - Problem 1.43.

The specific heat capacity of water is  $1 \text{ cal g}^{-1} \text{ }^\circ\text{C}^{-1} = 4.186 \text{ J g}^{-1} \text{ }^\circ\text{C}^{-1}$ , so how does this translate into the heat capacity per water molecule? The molar weight of water is  $18.01528 \text{ g mol}^{-1}$  so in 1 gram of water there are

$$(0.1) \quad N = \frac{6.02 \times 10^{23}}{18.01528} = 3.3416 \times 10^{22} \text{ molecules}$$

The heat capacity of a single molecule is therefore

$$(0.2) \quad C = \frac{4.186}{3.3416 \times 10^{22}} = 1.2527 \times 10^{-22} \text{ J }^\circ\text{C}^{-1}$$

$$(0.3) \quad = 9.08k$$

where  $k$  is Boltzmann's constant.

Thus to raise the temperature of a sample of water by  $1^\circ \text{C}$ , we need to raise the energy of each molecule by around  $9k$ . If we assumed that all the thermal energy of water is due to quadratic degrees of freedom, then because each degree of freedom contributes  $\frac{1}{2}kT$  to the energy, we would need 18 degrees of freedom in a water molecule. As we've seen, water has around 10 such degrees of freedom, so the thermal energy must also be stored elsewhere.