

THIRD LAW OF THERMODYNAMICS; RESIDUAL ENTROPY

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Daniel V. Schroeder, *An Introduction to Thermal Physics*, (Addison-Wesley, 2000) - Problem 3.9.

The entropy is related to temperature by

$$\frac{1}{T} = \left(\frac{\partial S}{\partial U} \right)_{N,V} \quad (1)$$

Using the chain rule, and keeping everything at constant N and V , we can measure the change in entropy due to a change in temperature as

$$dS = \frac{dU}{T} = \left(\frac{\partial U}{\partial T} \right)_{N,V} \frac{dT}{T} = C_V \frac{dT}{T} \quad (2)$$

where C_V is the heat capacity at constant volume:

$$C_V = \left(\frac{\partial U}{\partial T} \right)_{N,V} \quad (3)$$

If we know $C_V(T)$ as a function of temperature, we can therefore find the change in entropy for a finite change in temperature by integration:

$$\Delta S = S_f - S_i = \int_{T_i}^{T_f} \frac{C_V(T)}{T} dT \quad (4)$$

The total entropy in a system at temperature T_f could theoretically be found by setting $T_i = 0$ in the integral

$$S_f - S(0) = \int_0^{T_f} \frac{C_V(T)}{T} dT \quad (5)$$

In theory, at absolute zero, any system should be in its (presumably) unique lowest energy state so the multiplicity of the zero state is 1, meaning that $S(0) = 0$, and this integral does in fact give the actual entropy in a system at temperature T_f . It's also obvious that for this integral to be finite (and positive) $C_V \rightarrow 0$ as $T \rightarrow 0$ at a rate such that the integral doesn't diverge at its lower limit. Thus we must have $C_V(T) \propto T^a$ where $a > 0$ as $T \rightarrow 0$. Either of these conditions is a statement of the *third law of thermodynamics*, which basically says that at absolute zero, the entropy of any system is zero.

In practice, as a substance is cooled, its molecular configuration can get frozen into one of several possible ground states, so that there is a *residual entropy* even when $T = 0$ K.

Example. Carbon monoxide molecules are linear and in the solid form, they can line up in two orientations: OC and CO. Thus at absolute zero, the collection of molecules can be considered as a frozen-in matrix of molecules oriented randomly, so for a sample of N molecules, there are 2^N possible structures. For a mole, the residual entropy is therefore

$$S_{res} = k \ln 2^{6.02 \times 10^{23}} = (1.38 \times 10^{-23}) (6.02 \times 10^{23}) \ln 2 = 5.76 \text{ J K}^{-1} \quad (6)$$

PINGBACKS

Pingback: Entropy changes in macroscopic systems

Pingback: Entropy of aluminum at low temperatures

Pingback: Entropy of a star

Pingback: Heat capacities using Maxwell relations

Pingback: Phases of helium-3

Pingback: Calcium carbonate phase diagram

Pingback: Diamond-graphite phase boundary