

ENTROPY OF MIXING IN A SMALL SYSTEM

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).


Reference: Daniel V. Schroeder, *An Introduction to Thermal Physics*, (Addison-Wesley, 2000) - Problem 5.57.

As an example of the entropy changes when two pure substances are mixed, consider a system of 100 molecules, which may vary in composition from 100% of species *A* through a mixture of *A* and *B* to 100% pure *B*. The entropy of mixing is given by

$$\Delta S_{\text{mixing}} = -Nk[x \ln x + (1-x) \ln(1-x)]$$

where $N = N_A + N_B$ is the fixed total number of molecules (100 here) and $x = N_A/N$.

For a small system such as this, we can generate an array of $\Delta S_{\text{mixing}}/k$ values for each value of N_A from 0 to 100. Plotting this as a bar chart, we get



Starting from $N_A = 0$ where $\Delta S/k = 0$ (since there is only one species at this point, there is no mixing), we see that the entropy increase per molecule as we convert successive molecules from A to B decreases. The changes in $\Delta S/k$ for the first few steps are:

add molecule number	change in $\frac{\Delta S}{k}$
1	5.60
2	4.20
3	3.67
4	3.32
5	3.06

The rate at which the entropy increases declines as we convert more molecules from A to B . If we add a slight impurity into an initially pure mixture of 100% B , this generates a larger increase in entropy than if we add a bit more impurity to an already mixed system.