

DIAGONALIZATION OF MATRICES

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: edX online course MIT 8.05 Week 6.

Suppose we have an operator T that has a matrix representation $T(\{v\})$ in some basis v . In some cases (not all!) it is possible to transform to a different basis u in which $T(\{u\})$ is a diagonal matrix. If the operator A transforms from the basis v to u , then we've seen that T transforms according to

$$(0.1) \quad T(\{u\}) = A^{-1}T(\{v\})A$$

The diagonalization problem is therefore to find the matrix A (if it exists). Assuming A does exist, we can look at the situation in two ways.

- (1) The matrix representation of T is diagonal in the u basis.
- (2) The operator $A^{-1}TA$ is diagonal in the original v basis.

To prove (2), we start with the fact that T is diagonal in the u basis, so that (no implied sums in what follows):

$$(0.2) \quad Tu_i = \lambda_i u_i$$

$$(0.3) \quad TAv_i = \lambda_i Av_i$$

$$(0.4) \quad A^{-1}TA v_i = \lambda_i A^{-1}Av_i$$

$$(0.5) \quad = \lambda_i v_i$$

From the last line, we see that v_i is an eigenvector of the operator $A^{-1}TA$ with the same eigenvalue λ_i as the eigenvector u_i of the operator T .

Thus we can write, in the original basis v , the equation

$$(0.6) \quad D_T \equiv A^{-1}TA$$

where D_T is the diagonalized version of the matrix T . This is known as a *similarity transformation*.

All this is fine, but we still haven't seen how to find A . It turns out that the columns of A are the eigenvectors of T . We can see this from the following argument.

In the basis v , each v_i has the form of a column vector with a 1 in the i th position and zeroes everywhere else. Thus the transformation to the u basis can be written as

$$(0.7) \quad u_k = Av_k = \begin{bmatrix} A_{11} & \dots & A_{1n} \\ \vdots & \ddots & \vdots \\ A_{n1} & \dots & A_{nn} \end{bmatrix} \begin{bmatrix} \vdots \\ 1 \\ \vdots \end{bmatrix} = \begin{bmatrix} A_{1k} \\ \vdots \\ A_{nk} \end{bmatrix}$$

The 1 is in the k th position in the column vector representing v_k and picks out the elements in column k of A in the product Av_k . Since u_k is the k th eigenvector of T , the columns of A are the eigenvectors of T .

Example. Suppose we have

$$(0.8) \quad T = \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix}$$

The eigenvalues are found in the usual way, from the characteristic determinant, which is

$$(0.9) \quad (1 - \lambda)^2 - 6 = 0$$

$$(0.10) \quad \lambda = 1 \pm \sqrt{6}$$

We can find the eigenvectors by solving $Tu_i = \lambda_i u_i$ for each eigenvalue, where u_i is a 2-element column vector. (I won't go through this, since it's just algebra.) Placing the eigenvectors as the columns in a matrix A , we have

$$(0.11) \quad A = \begin{bmatrix} \frac{\sqrt{6}}{3} & -\frac{\sqrt{6}}{3} \\ 1 & 1 \end{bmatrix}$$

The inverse of a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is

$$(0.12) \quad A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

So for our matrix, we have

$$(0.13) \quad A^{-1} = \begin{bmatrix} \frac{\sqrt{6}}{4} & \frac{1}{2} \\ -\frac{\sqrt{6}}{4} & \frac{1}{2} \end{bmatrix}$$

Doing the matrix products (just a lot of arithmetic) we get

$$(0.14) \quad A^{-1}TA = \begin{bmatrix} 1 + \sqrt{6} & 0 \\ 0 & 1 - \sqrt{6} \end{bmatrix}$$

The resulting matrix is diagonal, and the diagonal entries are the eigenvalues of T . It's worth noticing that the traces of $A^{-1}TA$ and T are the same (both = 2), as are the determinants (both = -5).

PINGBACKS

Pingback: Normal operators

Pingback: Spectral theorem for normal operators

Pingback: Eigenvalues and eigenvectors - examples

Pingback: Adding two spin-1/2 systems - product and total-s bases

Pingback: Total J for sum of two angular momenta