

## HERMITIAN OPERATORS - A FEW EXAMPLES

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Exercise 1.6.2.

Here are a few more results about hermitian operators.

Suppose we are given two hermitian operators  $\Omega$  and  $\Lambda$ . We'll look at some combinations of these operators.

The operator  $\Omega\Lambda$  has the hermitian conjugate

$$(0.1) \quad (\Omega\Lambda)^\dagger = \Lambda^\dagger\Omega^\dagger = \Lambda\Omega$$

Thus the product operator  $\Omega\Lambda$  is hermitian only if  $\Lambda$  and  $\Omega$  commute.

The operator  $\Omega\Lambda + \lambda\Omega$  for some complex scalar  $\lambda$  has the hermitian conjugate

$$(0.2) \quad (\Omega\Lambda + \lambda\Omega)^\dagger = \Lambda^\dagger\Omega^\dagger + \lambda^*\Omega^\dagger$$

$$(0.3) \quad = \Lambda\Omega + \lambda^*\Omega$$

This operator is therefore hermitian only if  $\Lambda$  and  $\Omega$  commute and  $\lambda$  is real.

The commutator has the hermitian conjugate

$$(0.4) \quad [\Omega, \Lambda]^\dagger = (\Omega\Lambda - \Lambda\Omega)^\dagger$$

$$(0.5) \quad = \Lambda\Omega - \Omega\Lambda$$

$$(0.6) \quad = [\Lambda, \Omega]$$

$$(0.7) \quad = -[\Omega, \Lambda]$$

Thus the commutator is anti-hermitian (the hermitian conjugate is the negative of the original operator).

Finally, what happens if we multiply the commutator by  $i$ ?

$$(0.8) \quad (i[\Omega, \Lambda])^\dagger = -i(\Omega\Lambda - \Lambda\Omega)^\dagger$$

$$(0.9) \quad = -i(\Lambda\Omega - \Omega\Lambda)$$

$$(0.10) \quad = -i[\Lambda, \Omega]$$

$$(0.11) \quad = i[\Omega, \Lambda]$$

Thus the operator  $i[\Omega, \Lambda]$  is hermitian.