

HAMILTONIAN FOR THE TWO-BODY PROBLEM

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Section 2.5; Exercise 2.5.4.

Here we derive the equations of motion of the two-body problem using the Hamiltonian formalism.

The Hamiltonian is given by

$$(0.1) \quad H(q, p) = \sum_i p_i \dot{q}_i - L(q, \dot{q})$$

where the velocities \dot{q}_i are expressed in terms of the positions q_i and momenta p_i . In this case, we start with the Lagrangian in terms of the centre of mass position \mathbf{r}_{CM} and the relative position \mathbf{r} of mass 2 to mass 1.

$$(0.2) \quad L = \frac{1}{2} (m_1 + m_2) |\dot{\mathbf{r}}_{CM}|^2 + \frac{1}{2} \frac{m_1 m_2}{m_1 + m_2} |\dot{\mathbf{r}}|^2 - V(\mathbf{r})$$

$$(0.3) \quad = \frac{M}{2} |\dot{\mathbf{r}}_{CM}|^2 + \frac{\mu}{2} |\dot{\mathbf{r}}|^2 - V(\mathbf{r})$$

where $M = m_1 + m_2$ is the total mass and $\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass.

There are potentially 6 velocity components and 6 coordinate components in the Lagrangian, but the 3 components of \mathbf{r}_{CM} do not appear, which simplifies things a bit. To convert to a Hamiltonian, we need the momenta

$$(0.4) \quad p_i = \frac{\partial L}{\partial \dot{q}_i}$$

The x component of momentum of the centre of mass is

$$(0.5) \quad p_{CM,x} = \frac{\partial L}{\partial \dot{r}_{CM,x}} = M \dot{r}_{CM,x}$$

The other two components of the centre of mass velocity, and of the relative velocity, have a similar form, and in general we can write

$$(0.6) \quad p_{CM,i} = M \dot{r}_{CM,i}$$

$$(0.7) \quad p_i = \mu \dot{r}_i$$

In vector notation, this becomes

$$(0.8) \quad \dot{\mathbf{r}}_{CM} = \frac{\mathbf{p}_{CM}}{M}$$

$$(0.9) \quad \dot{\mathbf{r}} = \frac{\mathbf{p}}{\mu}$$

$$(0.10) \quad |\dot{\mathbf{r}}_{CM}|^2 = \frac{|\mathbf{p}_{CM}|^2}{M^2}$$

$$(0.11) \quad |\dot{\mathbf{r}}|^2 = \frac{|\mathbf{p}|^2}{\mu^2}$$

The Lagrangian thus becomes

$$(0.12) \quad L = \frac{|\mathbf{p}_{CM}|^2}{2M} + \frac{|\mathbf{p}|^2}{2\mu} - V(\mathbf{r})$$

The Hamiltonian is

$$(0.13) \quad H = \mathbf{p} \cdot \dot{\mathbf{r}} + \mathbf{p}_{CM} \cdot \dot{\mathbf{r}}_{CM} - L$$

$$(0.14) \quad = \frac{|\mathbf{p}|^2}{\mu} + \frac{|\mathbf{p}_{CM}|^2}{M} - \left[\frac{|\mathbf{p}_{CM}|^2}{2M} + \frac{|\mathbf{p}|^2}{2\mu} - V(\mathbf{r}) \right]$$

$$(0.15) \quad = \frac{|\mathbf{p}_{CM}|^2}{2M} + \frac{|\mathbf{p}|^2}{2\mu} + V(\mathbf{r})$$

Once we've got the Hamiltonian, we can apply Hamilton's canonical equations to get the equations of motion.

$$(0.16) \quad \frac{\partial H}{\partial p_i} = \dot{r}_i$$

$$(0.17) \quad -\frac{\partial H}{\partial r_i} = \dot{p}_i$$

Since \mathbf{r}_{CM} does not appear in the Hamiltonian, we have

$$(0.18) \quad \dot{\mathbf{p}}_{CM} = 0$$

$$(0.19) \quad \mathbf{p}_{CM} = \text{constant}$$

so the momentum of the centre of mass does not change, as expected.
For \mathbf{r} , we have

$$(0.20) \quad \frac{\partial H}{\partial p_i} = \frac{p_i}{\mu} = \dot{r}_i$$

$$(0.21) \quad \frac{\partial H}{\partial r_i} = \frac{\partial V}{\partial r_i} = -\dot{p}_i$$

The first equation tells us nothing new, while the second is just Newton's law for a central force: $\dot{\mathbf{p}} = -\nabla V$.

PINGBACKS

Pingback: Canonical transformations: a few more examples