

HAMILTONIAN FOR THE ELECTROMAGNETIC FORCE

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Section 2.6.

Here we derive the equations of motion for the electromagnetic force using the Hamiltonian formalism.

The Hamiltonian is given by

$$(0.1) \quad H(q, p) = \sum_i p_i \dot{q}_i - L(q, \dot{q})$$

where the velocities \dot{q}_i are expressed in terms of the positions q_i and momenta p_i . The electromagnetic Lagrangian is

$$(0.2) \quad L = \frac{1}{2} m \mathbf{v} \cdot \mathbf{v} - q\phi + \frac{q}{c} \mathbf{v} \cdot \mathbf{A}$$

where ϕ is the electric potential and \mathbf{A} is the magnetic potential, with \mathbf{v} the velocity of the charge q with mass m . To convert to the Hamiltonian, we need the momentum, defined as

$$p_i = \frac{\partial L}{\partial \dot{q}_i}$$

In this case, the generalized velocity is given by

$$(0.3) \quad \dot{q}_i = v_i$$

so we have

$$(0.4) \quad p_i = mv_i + \frac{q}{c} A_i$$

or, in vector notation

$$(0.5) \quad \mathbf{p} = m\mathbf{v} + \frac{q}{c} \mathbf{A}$$

$$(0.6) \quad \mathbf{v} = \frac{\mathbf{p}}{m} - \frac{q}{mc} \mathbf{A}$$

The Lagrangian is therefore

$$(0.7) \quad L = \frac{|\mathbf{p} - q\mathbf{A}/c|^2}{2m} - q\phi + \frac{q}{c} \left(\frac{\mathbf{p}}{m} - \frac{q}{mc} \mathbf{A} \right) \cdot \mathbf{A}$$

The first sum in the Hamiltonian is

$$(0.8) \quad \sum_i p_i \dot{q}_i = \mathbf{p} \cdot \mathbf{v} = \mathbf{p} \cdot \left(\frac{\mathbf{p}}{m} - \frac{q}{mc} \mathbf{A} \right)$$

The Hamiltonian is then

$$(0.9) \quad H = \mathbf{p} \cdot \left(\frac{\mathbf{p}}{m} - \frac{q}{mc} \mathbf{A} \right) - \frac{|\mathbf{p} - q\mathbf{A}/c|^2}{2m} + q\phi - \frac{q}{c} \left(\frac{\mathbf{p}}{m} - \frac{q}{mc} \mathbf{A} \right) \cdot \mathbf{A}$$

$$(0.10) \quad = \left(\frac{\mathbf{p}}{m} - \frac{q}{mc} \mathbf{A} \right) \cdot \left(\mathbf{p} - \frac{q}{c} \mathbf{A} \right) - \frac{|\mathbf{p} - q\mathbf{A}/c|^2}{2m} + q\phi$$

$$(0.11) \quad = \frac{|\mathbf{p} - q\mathbf{A}/c|^2}{2m} + q\phi$$

PINGBACKS

Pingback: Harmonic oscillator in a magnetic field

Pingback: Second-order correction to zeeman effect in hydrogen