

CORRESPONDENCE BETWEEN CLASSICAL AND QUANTUM TRANSFORMATIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Chapter 11.

When we consider infinitesimal transformations of some dynamical variable, there is a correspondence between classical and quantum mechanics which we can see as follows. First, we'll summarize the results from classical mechanics. We can define a canonical transformation generated by a variable g as

$$(0.1) \quad \bar{q}_i = q_i + \varepsilon \frac{\partial g}{\partial p_i} \equiv q_i + \delta q_i$$

$$(0.2) \quad \bar{p}_i = p_i - \varepsilon \frac{\partial g}{\partial q_i} \equiv p_i + \delta p_i$$

Here, ε is an infinitesimal amount and δq_i and δp_i are the infinitesimal amounts by which the coordinates and momenta vary. It follows from these definitions that, for any dynamical variable ω , its variation $\delta\omega$ is given by a Poisson bracket

$$(0.3) \quad \delta\omega = \omega(\bar{q}_i, \bar{p}_i) - \omega(q_i, p_i) = \varepsilon \{ \omega, g \}$$

For the special cases of coordinates and momenta, this is

$$(0.4) \quad \delta q_i = \varepsilon \{ q_i, g \}$$

$$(0.5) \quad \delta p_i = \varepsilon \{ p_i, g \}$$

If the generator is the momentum p_j , then

$$(0.6) \quad \delta q_i = \varepsilon \{ q_i, p_j \} = \varepsilon \delta_{ij}$$

$$(0.7) \quad \delta p_i = \varepsilon \{ p_i, p_j \} = 0$$

Thus, in classical mechanics, p_j is the generator of translations in direction j .

If $\omega = H$ (the Hamiltonian) and if $\{H, g\} = 0$, then g is conserved (it doesn't vary with time). Because the transformation 0.1 and 0.2 is canonical, it preserves the Poisson brackets so that

$$(0.8) \quad \{\bar{q}_i, \bar{q}_j\} = \{\bar{p}_i, \bar{p}_j\} = 0$$

$$(0.9) \quad \{\bar{q}_i, \bar{p}_j\} = \delta_{ij}$$

What do these things correspond to in quantum mechanics? [I find Shankar's treatment in section 11.2 to be almost tautological, since it merely repeats the derivation given earlier. I'll try to be a bit more general.]

Suppose we have some infinitesimal transformation given by a unitary operator $U(\varepsilon)$. We can then define the changes in X and P by

$$(0.10) \quad \delta X = U^\dagger(\varepsilon) X U(\varepsilon) - X$$

$$(0.11) \quad \delta P = U^\dagger(\varepsilon) P U(\varepsilon) - P$$

Since $U(\varepsilon)$ describes an infinitesimal transformation, we can expand it to first order in ε :

$$(0.12) \quad U(\varepsilon) = I - \frac{i\varepsilon}{\hbar} G$$

where $G = G^\dagger$ is some Hermitian operator known as the generator of the transformation. (We've seen a proof that the translation operator $T(\varepsilon)$ (a special case of $U(\varepsilon)$) is unitary and that its generator is Hermitian earlier, and the current case follows the same reasoning.) Using this form we have from 0.10 and 0.11, to order ε :

$$(0.13) \quad \delta X = \left(I + \frac{i\varepsilon}{\hbar} G \right) X \left(I - \frac{i\varepsilon}{\hbar} G \right) - X$$

$$(0.14) \quad = -\frac{i\varepsilon}{\hbar} [X, G]$$

$$(0.15) \quad \delta P = \left(I + \frac{i\varepsilon}{\hbar} G \right) P \left(I - \frac{i\varepsilon}{\hbar} G \right) - P$$

$$(0.16) \quad = -\frac{i\varepsilon}{\hbar} [P, G]$$

If $G = P$, then

$$(0.17) \quad \delta X = -\frac{i\varepsilon}{\hbar} [X, P] = \varepsilon I$$

$$(0.18) \quad \delta P = -\frac{i\varepsilon}{\hbar} [P, P] = 0$$

Comparing this with 0.6 and 0.7 we see that (in one dimension, where the classical coordinate is given by x and momentum by p) there is a correspondence between the classical Poisson bracket and quantum commutator:

$$(0.19) \quad \{x, p\} \leftrightarrow -\frac{i}{\hbar} [X, P]$$

The momentum operator P in quantum mechanics is thus the generator of translations, just as p generates translations in classical mechanics.

More generally, we can define the variation in some arbitrary dynamical operator Ω in a similar way, using 0.12 to expand the RHS:

$$(0.20) \quad \delta\Omega = U^\dagger(\varepsilon)\Omega U(\varepsilon) - \Omega$$

$$(0.21) \quad = -\frac{i\varepsilon}{\hbar} [\Omega, G]$$

The correspondence with classical mechanics is then

$$(0.22) \quad \{\omega, g\} \leftrightarrow -\frac{i}{\hbar} [\Omega, G]$$

The general rule is that a quantum commutator is $i\hbar$ times the corresponding classical Poisson bracket.

If $\Omega = H$ and $[H, G] = 0$, then by Ehrenfest's theorem, $\langle \dot{G} \rangle = 0$ and the average value of G is conserved.

The correspondence is a bit odd in that the generator g in classical mechanics enters as a derivative in 0.1 and 0.2 while the generator G in quantum mechanics enters as an operator (no derivatives) in 0.12.

One other feature is worth noting. A canonical transformation preserves the Poisson brackets 0.8 in the new coordinate system. In quantum mechanics, it is the commutators that get preserved. For example, using the fact that U is unitary so that $UU^\dagger = I$:

$$(0.23) \quad U^\dagger [X, P] U = U^\dagger X P U - U^\dagger P X U$$

$$(0.24) \quad = U^\dagger X U U^\dagger P U - U^\dagger P U U^\dagger X U$$

$$(0.25) \quad = [U^\dagger X U, U^\dagger P U]$$

PINGBACKS

Pingback: [Finite transformations: correspondence between classical and quantum](#)

Pingback: [Harmonic oscillator in a magnetic field](#)