

## EIGENVALUES OF TWO-DIMENSIONAL ANGULAR MOMENTUM

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press.  
Chapter 12, Exercise 12.3.1.

[If some equations are too small to read easily, use your browser's magnifying option (Ctrl + on Chrome, probably something similar on other browsers).]

The angular momentum operator  $L_z$  for rotations in two dimensions has the form, in polar coordinates, of

$$(0.1) \quad L_z = -i\hbar \frac{\partial}{\partial \phi}$$

To find the eigenvalues and eigenfunctions, we need to solve

$$(0.2) \quad L_z |\ell_z\rangle = \ell_z |\ell_z\rangle$$

where  $|\ell_z\rangle$  is the eigenfunction and  $\ell_z$  is the corresponding eigenvalue. Using polar coordinates, we must solve

$$(0.3) \quad -i\hbar \frac{\partial}{\partial \phi} \psi_{\ell_z}(\rho, \phi) = \ell_z \psi_{\ell_z}(\rho, \phi)$$

where  $\rho$  is the radial coordinate. As the only derivative here is with respect to  $\phi$ , we can solve this using separation of variables by proposing a solution of form

$$(0.4) \quad \psi_{\ell_z}(\rho, \phi) = R(\rho) \Phi(\phi)$$

Substituting this and cancelling off  $R(\rho)$  we get

$$(0.5) \quad -i\hbar \frac{\partial}{\partial \phi} \Phi(\phi) = \ell_z \Phi(\phi)$$

which has the solution

$$(0.6) \quad \Phi(\phi) = A e^{i\ell_z \phi / \hbar}$$

for some constant  $A$ , which we can absorb into  $R(\rho)$  to give the general solution

$$(0.7) \quad \psi_{\ell_z}(\rho, \phi) = R(\rho) e^{i\ell_z \phi / \hbar}$$

[This is actually the two-dimensional version of the more general 3-d case, in which the solution involved a radial function multiplied by a spherical harmonic.]

At this stage, the eigenvalue  $\ell_z$  could be any number, real or complex, since they all satisfy **??**. However, since  $L_z$  is an observable, it must be hermitian, which implies that  $L_z^\dagger = L_z$ , so that

$$(0.8) \quad \langle \psi_1 | L_z | \psi_2 \rangle = \langle \psi_2 | L_z | \psi_1 \rangle^*$$

In the coordinate basis, we have

$$(0.9) \quad \int_0^\infty \int_0^{2\pi} \psi_1^* \left( -i\hbar \frac{\partial}{\partial \phi} \right) \psi_2 d\phi d\rho = \left[ \int_0^\infty \int_0^{2\pi} \psi_2^* \left( -i\hbar \frac{\partial}{\partial \phi} \right) \psi_1 d\phi d\rho \right]^*$$

Integrating the LHS by parts, we have

$$(0.10) \quad \int_0^\infty \int_0^{2\pi} \psi_1^* \left( -i\hbar \frac{\partial}{\partial \phi} \right) \psi_2 d\phi d\rho = -i\hbar \int_0^\infty \psi_1^* \psi_2 \Big|_0^{2\pi} d\rho + i\hbar \int_0^\infty \int_0^{2\pi} \frac{\partial \psi_1^*}{\partial \phi} \psi_2 d\phi d\rho$$

The second term on the RHS is seen to be equal to the RHS of **??**, so in order for **??** to be true, we must have

$$(0.11) \quad \int_0^\infty \psi_1^* \psi_2 \Big|_0^{2\pi} d\rho = 0$$

Although two different eigenfunctions  $\psi_1$  and  $\psi_2$  are orthogonal and thus would satisfy this condition automatically, the condition must also be true when  $\psi_1 = \psi_2$ . This gives us the condition that

$$(0.12) \quad \psi_{\ell_z}(2\pi) = \psi_{\ell_z}(0)$$

That is, the eigenfunctions must be periodic with period  $2\pi$ . Looking back at **??**, we see that this forces the eigenvalues  $\ell_z$  to be integral multiples of  $\hbar$ :

$$(0.13) \quad \ell_z = m\hbar$$

$$(0.14) \quad m = 0, \pm 1, \pm 2, \dots$$

Here  $m$  is the *magnetic quantum number*, not the mass.

#### PINGBACKS

Pingback: Eigenvalues of angular momentum

Pingback: Angular momentum: probabilities of eigenvalues in two dimensions

Pingback: Radially symmetric potentials, angular momentum and centrifugal force

Pingback: Angular momentum of circular motion

Pingback: Harmonic oscillator in 2 dimensions: comparison with rectangular coordinates