

## SPHERICALLY SYMMETRIC POTENTIALS: A SIMPLE EXAMPLE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Chapter 12, Exercise 12.6.1.

[If some equations are too small to read easily, use your browser's magnifying option (Ctrl + on Chrome, probably something similar on other browsers).]

The Schrödinger equation in 3-d for a potential that depends only on  $r$  is

$$(0.1) \quad -\frac{\hbar^2}{2\mu} \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \left( \frac{\partial^2 \psi}{\partial \phi^2} \right) \right] + V \psi = E \psi$$

The angular part of the operator on the LHS is essentially the angular momentum operator  $L^2$  (times  $1/2\mu r^2$ ):

$$(0.2) \quad L^2 = -\hbar^2 \left[ \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right]$$

, so we can write this as

$$(0.3) \quad -\frac{\hbar^2}{2\mu} \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \psi}{\partial r} \right) + V \psi + \frac{L^2}{2\mu r^2} \psi = E \psi$$

Eigenfunctions in this equation satisfy

$$(0.4) \quad \psi = R_{Elm}(r) Y_l^m(\theta, \phi)$$

where the subscript  $Elm$  refers to the energy  $E$  and the angular momentum quantum numbers  $l$  and  $m$ .  $Y_l^m$  is a spherical harmonic and  $R_{Elm}$  is the radial function which depends on the potential  $V$ . The eigenvalues of  $L^2$  are  $l(l+1)\hbar^2$  so 0.3 becomes

$$(0.5) \quad -\frac{\hbar^2}{2\mu} \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial R_{El}}{\partial r} \right) + \frac{l(l+1)\hbar^2}{2\mu r^2} R_{El} + V R_{El} = E R_{El}$$

We've dropped the  $m$  from  $R_{Elm}$  since, for a spherically symmetric potential, the radial function is independent of  $m$ .

**Example.** Suppose a particle is described by the wave function

$$(0.6) \quad \psi_E(r, \theta, \phi) = Ae^{-r/a_0}$$

where  $A$  and  $a_0$  are constants. What can we deduce about the system?

First, since  $\psi_E$  is independent of  $\theta$  and  $\phi$  we see from 0.2 that

$$(0.7) \quad L^2\psi_E = 0$$

so the eigenvalue is  $l = 0$  and the state has no angular momentum. From 0.3 we therefore have

$$(0.8) \quad -\frac{\hbar^2}{2\mu} \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \psi}{\partial r} \right) + V\psi = E\psi$$

Working out the derivatives, we have

$$(0.9) \quad \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \psi}{\partial r} \right) = -\frac{A}{r^2} \frac{d}{dr} \left( \frac{r^2}{a_0} e^{-r/a_0} \right)$$

$$(0.10) \quad = Ae^{-r/a_0} \left( -\frac{2}{ra_0} + \frac{1}{a_0^2} \right)$$

Plugging this back into 0.8 and cancelling terms gives

$$(0.11) \quad -\frac{2}{ra_0} + \frac{1}{a_0^2} = \frac{2\mu}{\hbar^2} (V - E)$$

If  $V(r) \rightarrow 0$  as  $r \rightarrow \infty$  we have, in this limit

$$(0.12) \quad E = -\frac{\hbar^2}{2\mu a_0^2}$$

The energy is constant at all values of  $r$  so we can now find  $V$  from 0.11

$$(0.13) \quad -\frac{2}{ra_0} + \frac{1}{a_0^2} = \frac{2\mu}{\hbar^2} \left( V(r) + \frac{\hbar^2}{2\mu a_0^2} \right)$$

$$(0.14) \quad V(r) = -\frac{\hbar^2}{\mu a_0 r}$$