

SPINOR IN OSCILLATING MAGNETIC FIELD - PART 1

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Chapter 14, Exercise 14.4.3, Part 1.

[If some equations are too small to read easily, use your browser's magnifying option (Ctrl + on Chrome, probably something similar on other browsers).]

Classically, if a magnetic moment μ is placed in a magnetic field that precesses about the z axis, the magnetic moment itself precesses. If the field is given as

$$(0.1) \quad \mathbf{B} = B \cos \omega t \hat{\mathbf{x}} - B \sin \omega t \hat{\mathbf{y}} + B_0 \hat{\mathbf{z}}$$

then in a frame that rotates with the same frequency as the field, the magnetic field appears to be constant with value

$$(0.2) \quad \mathbf{B}_r = B \hat{\mathbf{x}}_r + \left(B_0 - \frac{\omega}{\gamma} \right) \hat{\mathbf{z}}$$

where

$$(0.3) \quad \hat{\mathbf{x}}_r = \cos \omega t \hat{\mathbf{x}} - \sin \omega t \hat{\mathbf{y}}$$

is a unit vector along the x axis in the rotating frame. We now want to see how this result transfers into quantum mechanics.

We begin with the Schrödinger equation for the state $|\psi(t)\rangle$ in the lab (non-rotating) frame, which is, as usual

$$(0.4) \quad i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle = H |\psi(t)\rangle$$

We'll study the case where $|\psi(t)\rangle$ is a spin $\frac{1}{2}$ state, for which the Hamiltonian is

$$(0.5) \quad H = -\gamma \mathbf{S} \cdot \mathbf{B}$$

We can analyze the situation in the rotating frame by applying a unitary rotation operator to the lab state. That is

$$(0.6) \quad |\psi_r(t)\rangle = e^{-i\omega t S_z/\hbar} |\psi(t)\rangle = e^{-i\omega t \sigma_z/2} |\psi(t)\rangle$$

$$(0.7) \quad = \left[\cos \frac{\omega t}{2} I - i \sin \frac{\omega t}{2} \sigma_z \right] |\psi(t)\rangle$$

[It seems to me that this unitary operator is for a rotation by an angle ωt , and since the rotation of the field in 0.1 is given by a frequency $-\omega \hat{\mathbf{z}}$, we should really be using the rotation operator $e^{i\omega t \sigma_z/2}$. However if we do this (I tried) we get the wrong answer, so presumably the transformation 0.6 is correct.]

Our first goal is to find the Schrödinger equation for $|\psi_r(t)\rangle$, which involves finding the corresponding Hamiltonian. The Schrödinger equation is

$$(0.8) \quad i\hbar \frac{\partial}{\partial t} |\psi_r(t)\rangle = H_r |\psi_r(t)\rangle$$

Inserting 0.6 into the LHS and differentiating, we get

$$(0.9) \quad i\hbar \frac{\partial}{\partial t} |\psi_r(t)\rangle = \frac{\hbar\omega\sigma_z}{2} e^{-i\omega t \sigma_z/2} |\psi(t)\rangle + i\hbar e^{-i\omega t \sigma_z/2} \frac{\partial}{\partial t} |\psi(t)\rangle$$

$$(0.10) \quad = \frac{\hbar\omega\sigma_z}{2} |\psi_r(t)\rangle + e^{-i\omega t \sigma_z/2} H |\psi(t)\rangle$$

$$(0.11) \quad = \frac{\hbar\omega\sigma_z}{2} |\psi_r(t)\rangle - e^{-i\omega t \sigma_z/2} \gamma \mathbf{S} \cdot \mathbf{B} |\psi(t)\rangle$$

We would like the RHS to be in the form of the RHS of 0.8, but in the second term, the problem is that $e^{-i\omega t \sigma_z/2}$ does not commute with \mathbf{S} so we can't just swap the $e^{-i\omega t \sigma_z/2}$ and $\mathbf{S} \cdot \mathbf{B}$ factors. We need to multiply out the terms and see what simplifications we can do.

In what follows, it's easier to work with the Pauli matrices defined by

$$(0.12) \quad \mathbf{S} = \frac{\hbar}{2} \boldsymbol{\sigma}$$

We'll also need a few theorems involving σ_i

$$(0.13) \quad \sigma_i \sigma_j = -\sigma_j \sigma_i$$

$$(0.14) \quad \sigma_i \sigma_j = \delta_{ij} I + i \sum_k \epsilon_{ijk} \sigma_k$$

We'll also define some shorthand for the trig functions:

$$(0.15) \quad c \equiv \cos \frac{\omega t}{2}$$

$$(0.16) \quad s \equiv \sin \frac{\omega t}{2}$$

$$(0.17) \quad c_1 \equiv \cos \omega t$$

$$(0.18) \quad s_1 \equiv \sin \omega t$$

The standard double-angle formulas are

$$(0.19) \quad c_1 = c^2 - s^2$$

$$(0.20) \quad s_1 = 2sc$$

Using 0.1 and 0.7 we have

$$(0.21) \quad i\omega t \sigma_z / 2 \gamma \mathbf{S} \cdot \mathbf{B} = -\frac{\gamma \hbar}{2} [B(c - is\sigma_z)(\sigma_x c_1 - \sigma_y s_1) + B_0(c - is\sigma_z)\sigma_z]$$

The last term on the RHS is in the correct form since there are no commutation problems here. So we need to work on the first term, which we'll isolate here:

$$(0.22) \quad (c - is\sigma_z)(\sigma_x c_1 - \sigma_y s_1) = c_1 c \sigma_x + ic_1 s \sigma_x \sigma_z - s_1 c \sigma_y - is_1 s \sigma_y \sigma_z$$

We can now use the identities 0.13 and 0.14 and the trig identities above to get

$$(0.23) \quad (c - is\sigma_z)(\sigma_x c_1 - \sigma_y s_1) = (c^2 - s^2) c \sigma_x + i(c^2 - s^2) s \sigma_x \sigma_z - 2sc^2 \sigma_y - 2is^2 c \sigma_y \sigma_z$$

$$(0.24) \quad = (c^2 - s^2) c \sigma_x + i(c^2 - s^2) s \sigma_x \sigma_z - 2isc^2 \sigma_x \sigma_z + 2s^2 c \sigma_x$$

$$(0.25) \quad = (c^3 - s^2 c + 2s^2 c) \sigma_x + i(-s^3 + c^2 s - 2sc^2) \sigma_x \sigma_z$$

$$(0.26) \quad = (c^2 + s^2) c \sigma_x - i(c^2 + s^2) s \sigma_x \sigma_z$$

$$(0.27) \quad = \sigma_x (c - is\sigma_z)$$

$$(0.28) \quad = \sigma_x e^{-i\omega t \sigma_z / 2}$$

Plugging this into 0.21 and then back into 0.11 we get

$$(0.29) \quad \frac{\partial}{\partial t} |\psi_r(t)\rangle = \frac{\hbar \omega \sigma_z}{2} |\psi_r(t)\rangle - \frac{\gamma \hbar}{2} [B\sigma_x + B_0\sigma_z] e^{-i\omega t \sigma_z / 2} |\psi(t)\rangle$$

$$(0.30) \quad = [(\omega - \gamma B_0) S_z - \gamma B S_x] |\psi_r(t)\rangle$$

Comparing this with 0.2, we see that we can write the result as

$$(0.31) \quad i\hbar \frac{\partial}{\partial t} |\psi_r(t)\rangle = -\gamma \mathbf{S} \cdot \mathbf{B}_r |\psi_r(t)\rangle$$

Thus in the rotating frame, the Schrödinger equation has the same form as the classical relation, with a time-independent magnetic field \mathbf{B}_r .

PINGBACKS

Pingback: Spinor in oscillating magnetic field - part 2