

SPIN FLIP OF ELECTRON IN MAGNETIC FIELD

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Chapter 14, Exercise 14.4.4.

[If some equations are too small to read easily, use your browser's magnifying option (Ctrl + on Chrome, probably something similar on other browsers).]

The fact that the rotation operator in 2-d spin space can be written in terms of the Pauli matrices as

$$(1) \quad U[R(\theta)] = e^{-i\theta \cdot \sigma / 2} = \cos \frac{\theta}{2} I - i \sin \frac{\theta}{2} (\hat{\theta} \cdot \sigma)$$

allows us to do some calculations involving an electron in a magnetic field. We've seen that placing a magnetic moment that is due to angular momentum in a constant magnetic field causes the magnetic moment to precess about the direction of the field. The Hamiltonian of an electron with spin \mathbf{S} in a constant field \mathbf{B} is

$$(2) \quad H = -\gamma \mathbf{S} \cdot \mathbf{B}$$
$$(3) \quad = -\frac{\gamma \hbar}{2} \boldsymbol{\sigma} \cdot \mathbf{B}$$

where γ is the gyromagnetic ratio

$$(4) \quad \gamma = \frac{-e}{m} \text{ (SI)} = \frac{-e}{mc} \text{ (CGS)}$$
$$(5) \quad = 1.76 \times 10^{11} \text{ s}^{-1} \text{T}^{-1}$$
$$(6) \quad = 1.76 \times 10^7 \text{ s}^{-1} \text{G}^{-1}$$

As an example, suppose we have an electron initially in the spin-up state, with $s_z = +\frac{\hbar}{2}$ and turn on a magnetic field of $\mathbf{B} = (100 \text{ G}) \hat{\mathbf{x}}$ at $t = 0$. As the applied field is perpendicular to the initial spin, the precession will cause the spin vector to rotate in the yz plane about the x axis. To find how long it takes the spin to flip, we need the propagator, which is

$$\begin{aligned}
 (7) \quad U(t) &= e^{-iHt/\hbar} \\
 (8) \quad &= e^{i\gamma\sigma \cdot \mathbf{B}t/2} \\
 (9) \quad &= e^{i\gamma Bt\sigma_x/2}
 \end{aligned}$$

Comparing with 1 we see that $U(t)$ is equivalent to a rotation operator with angle $\theta = -\gamma Bt$. Thus we have

$$\begin{aligned}
 (10) \quad U(t) &= \cos \frac{\gamma Bt}{2} I + i \sin \frac{\gamma Bt}{2} \sigma_x \\
 (11) \quad &= \begin{bmatrix} \cos \frac{\gamma Bt}{2} & i \sin \frac{\gamma Bt}{2} \\ i \sin \frac{\gamma Bt}{2} & \cos \frac{\gamma Bt}{2} \end{bmatrix}
 \end{aligned}$$

The electron's state as a function of time is thus

$$\begin{aligned}
 (12) \quad |\psi(t)\rangle &= U(t)|\psi(0)\rangle \\
 (13) \quad &= \begin{bmatrix} \cos \frac{\gamma Bt}{2} & i \sin \frac{\gamma Bt}{2} \\ i \sin \frac{\gamma Bt}{2} & \cos \frac{\gamma Bt}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \\
 (14) \quad &= \begin{bmatrix} \cos \frac{\gamma Bt}{2} \\ i \sin \frac{\gamma Bt}{2} \end{bmatrix}
 \end{aligned}$$

The state first has a 100% probability of being found with spin down when (using 6):

$$\begin{aligned}
 (15) \quad \sin \frac{\gamma Bt}{2} &= 1 \\
 (16) \quad t &= \frac{\pi}{\gamma B} \\
 (17) \quad &= \frac{\pi}{(1.76 \times 10^7)(100)} \\
 (18) \quad &= 1.78 \times 10^{-9} \text{ s}
 \end{aligned}$$