

## SPIN FLIP OF ELECTRON IN MAGNETIC FIELD

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press. Chapter 14, Exercise 14.4.4.

[If some equations are too small to read easily, use your browser's magnifying option (Ctrl + on Chrome, probably something similar on other browsers).]

The fact that the rotation operator in 2-d spin space can be written in terms of the Pauli matrices as

$$(0.1) \quad U[R(\theta)] = e^{-i\theta \cdot \sigma / 2} = \cos \frac{\theta}{2} I - i \sin \frac{\theta}{2} (\hat{\theta} \cdot \sigma)$$

allows us to do some calculations involving an electron in a magnetic field. We've seen that placing a magnetic moment that is due to angular momentum in a constant magnetic field causes the magnetic moment to precess about the direction of the field. The Hamiltonian of an electron with spin  $\mathbf{S}$  in a constant field  $\mathbf{B}$  is

$$(0.2) \quad H = -\gamma \mathbf{S} \cdot \mathbf{B}$$
$$(0.3) \quad = -\frac{\gamma \hbar}{2} \boldsymbol{\sigma} \cdot \mathbf{B}$$

where  $\gamma$  is the gyromagnetic ratio

$$(0.4) \quad \gamma = \frac{-e}{m} \text{ (SI)} = \frac{-e}{mc} \text{ (CGS)}$$

$$(0.5) \quad = 1.76 \times 10^{11} \text{ s}^{-1} \text{T}^{-1}$$

$$(0.6) \quad = 1.76 \times 10^7 \text{ s}^{-1} \text{G}^{-1}$$

As an example, suppose we have an electron initially in the spin-up state, with  $s_z = +\frac{\hbar}{2}$  and turn on a magnetic field of  $\mathbf{B} = (100 \text{ G}) \hat{\mathbf{x}}$  at  $t = 0$ . As the applied field is perpendicular to the initial spin, the precession will cause the spin vector to rotate in the  $yz$  plane about the  $x$  axis. To find how long it takes the spin to flip, we need the propagator, which is

$$(0.7) \quad U(t) = e^{-iHt/\hbar}$$

$$(0.8) \quad = e^{i\gamma\sigma \cdot \mathbf{B}t/2}$$

$$(0.9) \quad = e^{i\gamma Bt\sigma_x/2}$$

Comparing with 0.1 we see that  $U(t)$  is equivalent to a rotation operator with angle  $\theta = -\gamma Bt$ . Thus we have

$$(0.10) \quad U(t) = \cos \frac{\gamma Bt}{2} I + i \sin \frac{\gamma Bt}{2} \sigma_x$$

$$(0.11) \quad = \begin{bmatrix} \cos \frac{\gamma Bt}{2} & i \sin \frac{\gamma Bt}{2} \\ i \sin \frac{\gamma Bt}{2} & \cos \frac{\gamma Bt}{2} \end{bmatrix}$$

The electron's state as a function of time is thus

$$(0.12) \quad |\psi(t)\rangle = U(t) |\psi(0)\rangle$$

$$(0.13) \quad = \begin{bmatrix} \cos \frac{\gamma Bt}{2} & i \sin \frac{\gamma Bt}{2} \\ i \sin \frac{\gamma Bt}{2} & \cos \frac{\gamma Bt}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$(0.14) \quad = \begin{bmatrix} \cos \frac{\gamma Bt}{2} \\ i \sin \frac{\gamma Bt}{2} \end{bmatrix}$$

The state first has a 100% probability of being found with spin down when (using 0.6):

$$(0.15) \quad \sin \frac{\gamma Bt}{2} = 1$$

$$(0.16) \quad t = \frac{\pi}{\gamma B}$$

$$(0.17) \quad = \frac{\pi}{(1.76 \times 10^7)(100)}$$

$$(0.18) \quad = 1.78 \times 10^{-9} \text{ s}$$